

COMPUTING THE PARTITION FUNCTION FOR PERFECT MATCHINGS IN A HYPERGRAPH

ALEXANDER BARVINOK AND ALEX SAMORODNITSKY

November 2010

ABSTRACT. Given non-negative weights w_S on the k -subsets S of a km -element set V , we consider the sum of the products $w_{S_1} \cdots w_{S_m}$ over all partitions $V = S_1 \cup \dots \cup S_m$ into pairwise disjoint k -subsets S_i . When the weights w_S are positive and within a constant factor, fixed in advance, of each other, we present a simple polynomial time algorithm to approximate the sum within a polynomial in m factor. In the process, we obtain higher-dimensional versions of the van der Waerden and Bregman-Minc bounds for permanents. We also discuss applications to counting of perfect and nearly perfect matchings in hypergraphs.

1. INTRODUCTION AND MAIN RESULTS

Let us fix an integer $k > 1$. A collection $H \subset \binom{V}{k}$ of k -subsets of a finite set V is called a *k-uniform hypergraph* with *vertex* set V , while sets $S \in H$ are called *edges* of H . In particular, a uniform 2-hypergraph is an ordinary undirected graph on V without loops or multiple edges. A set $\{S_1, \dots, S_m\}$ of pairwise vertex disjoint edges of H such that $V = S_1 \cup \dots \cup S_m$ is called a *perfect matching* of hypergraph H . More generally, a *matching of size n* is a collection of n pairwise disjoint edges of H .

If a perfect matching exists then the number $|V|$ of vertices of V is divisible by k , so we have $|V| = km$ for some integer m . The hypergraph consisting of all k -subsets of V is called the *complete k-uniform hypergraph* with vertex set V . We denote it by $\binom{V}{k}$. A hypergraph is called a *complete k-partite hypergraph* if the set V of vertices is a union $V = V_1 \cup \dots \cup V_k$ of pairwise disjoint sets V_i , called *parts*, such that $|V_1| = \dots = |V_k| = m$ and the edges of the hypergraph are the subsets $S \subset V$ containing exactly one vertex in each part: $|S \cap V_1| = \dots = |S \cap V_k| = 1$. We denote such a hypergraph by $V_1 \times \dots \times V_k$.

Key words and phrases. hypergraph, perfect matching, partition function, permanent, van der Waerden inequality.

The research of the first author was partially supported by NSF Grant DMS 0856640. The research of the second author was partially supported by ISF grant 039-7165. The research of the authors was also partially supported by a United States - Israel BSF grant 2006377.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

We introduce the main object of the paper.

(1.1) Partition function. Let H be a k -uniform hypergraph with the set V of vertices such that $|V| = km$ for some positive integer m . Suppose that to every edge $S \in H$ a non-negative real number w_S is assigned. Such an assignment $W = \{w_S\}$ we call a *weight* on H . We say that W is *positive* if $w_S > 0$ for all $S \in H$. The polynomial

$$P_H(W) = \sum w_{S_1} \cdots w_{S_m},$$

where the sum is taken over all perfect matchings $\{S_1, \dots, S_m\}$ of H , is called the *partition function* of perfect matchings in hypergraph H . Sometimes we write just $P(W)$ if the choice of the hypergraph H is clear from the context.

We note that we can obtain the partition function $P_H(W)$ of an arbitrary k -uniform hypergraph $H \subset \binom{V}{k}$ by specializing $w_S = 0$ for $S \notin H$ in the partition function of the complete k -uniform hypergraph $\binom{V}{k}$.

The partition function of $\binom{V}{2}$ with $|V| = 2m$ is known as the *hafnian* of the $2m \times 2m$ symmetric matrix $A = (a_{ij})$, where a_{ij} is the weight of the edge consisting of the i -th and j -th vertices of V (diagonal elements of A can be chosen arbitrarily), see, for example, Section 8.2 of [Mi78]. If $V_1 \times V_2$ is a complete bipartite graph with $|V_1| = |V_2| = m$ then the corresponding partition function is the *permanent* of the $m \times m$ matrix $B = (b_{ij})$, where b_{ij} is the weight of the edge consisting of the i -th vertex of V_1 and j -th vertex of V_2 . The partition function of the complete k -partite hypergraph gives rise to a version of the permanent of a k -dimensional tensor, see, for example, [D87b].

In this paper, we address the problem of computing or approximating $P_H(W)$ efficiently. First, we define certain classes of weights W .

(1.2) Balanced and k -stochastic weights. We say that a positive weight $W = \{w_S\}$ on a k -uniform hypergraph is α -*balanced* for some $\alpha \geq 1$ if

$$\frac{w_{S_1}}{w_{S_2}} \leq \alpha \quad \text{for all } S_1, S_2 \in H.$$

Note that an α -balanced weight is also β -balanced for any $\beta > \alpha$.

Weight $Z = \{z_S\}$ is called *k -stochastic*, if

$$\sum_{\substack{S \in H \\ S \ni v}} z_S = 1 \quad \text{for all } v \in V.$$

In words: for every vertex, the sum of the weights of the edges containing the vertex is 1.

Now we are ready to state our first main result.

(1.3) Theorem. *Let us fix an integer $k > 1$ and a real $\alpha \geq 1$. Then there exists a real $\gamma = \gamma(k, \alpha) > 0$ such that if H is a complete k -uniform hypergraph or a complete k -partite hypergraph with km vertices and Z is a k -stochastic α -balanced weight on H then*

$$m^{-\gamma} e^{-m(k-1)} \leq P_H(Z) \leq m^{\gamma} e^{-m(k-1)}$$

provided $m > 1$.

In other words, for fixed k and α , the value of the partition function for a k -stochastic α -balanced weight on a complete k -uniform hypergraph or a complete k -partite hypergraph can vary only within a polynomial in m range.

The van der Waerden conjecture on permanents proved by Falikman [Fa81] and Egorychev [Eg81], see also [Gu08] for important new developments, asserts that if $A = (a_{ij})$ is an $n \times n$ doubly stochastic matrix, that is, a non-negative matrix with all row and column sums equal 1, then

$$\text{per } A \geq \frac{n!}{n^n} = \sqrt{2\pi n} e^{-n} \left(1 + O\left(\frac{1}{n}\right) \right).$$

A conjecture by Minc proved by Bregman [Br73], see also [Sc78] for a simpler proof, asserts that if $B = (b_{ij})$ is an $n \times n$ matrix with $b_{ij} \in \{0, 1\}$ for all i, j then

$$\text{per } B \leq \prod_{i=1}^n (r_i!)^{1/r_i},$$

where r_i is the i -th row sum of B . From this inequality one can deduce that if A is an $n \times n$ non-negative matrix with all row sums equal 1 and all the entries not exceeding α/n for some $\alpha \geq 1$ then

$$\text{per } A \leq n^{\gamma} e^{-n}$$

for some $\gamma = \gamma(\alpha) > 0$ and all $n > 1$, see [So03]. Thus the van der Waerden and Bregman-Minc inequalities together imply that $\text{per } A = e^{-n} n^{O(1)}$ for any $n \times n$ doubly stochastic matrix A whose entries are within a factor of $O(1)$ of each other. Theorem 1.3 presents an extension of this interesting fact to non-bipartite graphs for $k = 2$ and to hypergraphs for $k > 2$.

While there are generalizations of the Bregman-Minc bound to hafnians [AF08] and to higher-dimensional permanents [D87a], there are no straightforward generalizations of the van der Waerden bound, as the following two examples readily show.

Let $V = \{a_1, b_1, c_1, a_2, b_2, c_2\}$ be the set of vertices and let

$$\{a_1, b_1\}, \{b_1, c_1\}, \{a_1, c_1\}, \{a_2, b_2\}, \{b_2, c_2\}, \{a_2, c_2\}$$

be the set of edges of a graph G . Let us define a weight $Z = \{z_S\}$ on $\binom{V}{2}$ by choosing $z_S = 1/2$ if S is an edge of G and $z_S = 0$ otherwise. Then Z is a 2-stochastic weight on $\binom{V}{2}$ and $P(Z) = 0$. That is, the hafnian of a symmetric doubly stochastic matrix can be zero.

Let $V = \{a_1, a_2, b_1, b_2, c_1, c_2\}$ be the set of vertices and let

$$\{a_1, b_1, c_1\}, \{a_1, b_2, c_2\}, \{a_2, b_2, c_1\}, \{a_2, b_1, c_2\}$$

be the edges of a 3-partite hypergraph H . Let us define a weight $Z = \{z_S\}$ on $\{a_1, a_2\} \times \{b_1, b_2\} \times \{c_1, c_2\}$ by choosing $z_S = 1/2$ if $S \in H$ and $z_S = 0$ otherwise. Then Z is a 3-stochastic weight on $\{a_1, a_2\} \times \{b_1, b_2\} \times \{c_1, c_2\}$ and $P(Z) = 0$. Hence the permanent of a 3-stochastic 3-dimensional tensor can be zero.

Summarizing, for general k -stochastic weights Z there is no a priori non-zero lower bound for the partition function. If, however, we require Z to be α -balanced for any fixed $\alpha \geq 1$, the lower bound jumps to within a polynomial in m factor of the upper bound.

If H is a complete k -uniform hypergraph or a complete k -partite hypergraph, one can estimate $P_H(W)$ for any balanced but not necessarily k -stochastic weight W using scaling.

(1.4) Scaling. Let $W = \{w_S\}$ be a weight on the edges of a k -uniform hypergraph H with a vertex set V , where $|V| = km$. Let $\{\lambda_v > 0 : v \in V\}$ be reals. We say that a weight $Z = \{z_S\}$ on the hypergraph H is obtained from W by *scaling* if

$$z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all } S \in H.$$

It is easy to see that

$$P_H(Z) = \left(\prod_{v \in V} \lambda_v \right) P_H(W).$$

It turns out that any positive weight W on a complete k -uniform hypergraph or a complete k -partite hypergraph can be scaled to a k -stochastic weight Z (cf., for example, [Fr10] and Section 3 below). We show that the k -stochastic scaling of an α -balanced weight is α^{k+1} -balanced and obtain the following result.

(1.5) Theorem. *Let us fix an integer $k > 1$ and a real $\alpha \geq 1$. Then there exists a real $\gamma = \gamma(k, \alpha) > 0$ such that the following holds.*

Let H be a complete k -uniform hypergraph or a complete k -partite hypergraph with km vertices and let $W = \{w_S : S \in H\}$ be an α -balanced weight on H . Let us consider the function

$$f_W(X) = \sum_{S \in H} x_S \ln \frac{x_S}{w_S}$$

for a weight X on H . Let $\Omega_k(H)$ be the set of all k -stochastic weights on H and let

$$\zeta = \min_{X \in \Omega_k(H)} f_W(X).$$

Then

$$e^{-\zeta - m(k-1)} m^{-\gamma} \leq P_H(W) \leq e^{-\zeta - m(k-1)} m^\gamma.$$

The set $\Omega_k(H)$ is naturally identified with a convex polytope in \mathbb{R}^H . Function f is strictly convex and hence the optimization problem of computing ζ can be solved efficiently (in polynomial time) by interior point methods, see [NN94]. Thus Theorem 1.5 allows us to estimate the partition function of an α -balanced weight (for any $\alpha \geq 1$, fixed in advance) within a polynomial in m factor.

(1.6) A probabilistic interpretation. Let us fix $k > 1$ and let H be either a complete k -uniform hypergraph or a complete k -partite hypergraph with a set V of $|V| = km$ vertices. Let us fix $\alpha \geq 1$ and let $W = \{w_S : S \in H\}$ be an α -balanced weight on H . Let $|H|$ denote the number of edges of hypergraph H . Let us assume that

$$\sum_{S \in H} w_S = m,$$

in which case

$$\frac{m}{\alpha|H|} \leq w_S \leq \frac{\alpha m}{|H|} \quad \text{for all } S \in H.$$

In particular, for all sufficiently large m we have $w_S < 1$ for all $S \in H$, so we can introduce independent random Bernoulli variables X_S indexed by the edges $S \in H$, where

$$\Pr(X_S = 1) = w_S \quad \text{and} \quad \Pr(X_S = 0) = 1 - w_S.$$

For each vertex $v \in V$ let us define a random variable

$$Y_v = \sum_{\substack{S \in H \\ S \ni v}} X_S.$$

It is not hard show that

$$P_H(W) = \exp \left\{ m + O \left(\frac{1}{m^{k-2}} \right) \right\} \Pr \left(Y_v = 1 \quad \text{for all } v \in V \right).$$

For large m , the distribution of each random variable Y_v is approximately Poisson with

$$\mathbf{E} Y_v = \mu_v \quad \text{where} \quad \mu_v = \sum_{\substack{S \in H \\ S \ni v}} w_S,$$

so

$$\Pr(Y_v = 1) \approx \mu_v e^{-\mu_v}.$$

The probability of $Y_v = 1$ is maximized when $\mu_v = 1$, and when W is k -stochastic, the probabilities of $Y_v = 1$ are maximized simultaneously for all $v \in V$, so that

$$\Pr(Y_v = 1) \approx e^{-1} \quad \text{for all } v \in V.$$

Theorem 1.3 implies that in this case the events $Y_v = 1$ behave as if they were (almost) independent, so that

$$\Pr\left(Y_v = 1 \quad \text{for all } v \in V\right) \approx e^{-km}$$

up to a polynomial in m factor.

In Section 2, we discuss some combinatorial and algorithmic applications of Theorems 1.3 and 1.5. Namely, we present a simple polynomial time algorithm to distinguish hypergraphs having sufficiently many perfect matchings from hypergraphs that do not have nearly perfect matchings. We also prove a lower bound for the number of nearly perfect matchings in regular hypergraphs.

In the rest of the paper we prove Theorems 1.3 and 1.5.

In Section 3, we review some results about scaling. The results are not new, but we nevertheless provide proofs for completeness. In Section 4, we prove two crucial lemmas about scaling of α -balanced weights. In Section 5 we complete the proofs of Theorems 1.3 and 1.5.

Scaling was used in [L+00] to efficiently estimate permanents of non-negative matrices.

(1.7) Notation. As usual, for two functions f and g , where g is non-negative, we say that $f = O(g)$ if $|f| \leq \gamma g$ for some constant $\gamma > 0$. We will allow our constants γ to depend only on the dimension k of the hypergraph and the parameter $\alpha \geq 1$ in Definition 1.2 of an α -balanced weight.

2. COMBINATORIAL APPLICATIONS

Let us fix an integer $k > 1$ and let H be a k -uniform hypergraph with km vertices. As is known [Va79], the problem of counting perfect matchings in H is #P-hard. For $k = 2$ there is a classical polynomial time algorithm to check whether H has a perfect matching (see [LP09]) and a fully polynomial randomized approximation scheme is known for counting perfect matchings if H is bipartite [J+04]. For $k > 2$ finding if there is a perfect matching in H is an NP-complete problem even when H is k -partite [Ka72].

Theorem 1.5 allows us to distinguish in polynomial time between hypergraphs that have sufficiently many perfect matchings and hypergraphs that do not have nearly perfect matchings.

In this section, we let

$$\Phi_k(m) = \frac{(km)!}{(k!)^m m!}$$

be the number of perfect matchings in a complete k -uniform hypergraph with km vertices.

(2.1) Testing hypergraphs. Let us fix integer $k > 1$ and positive real $\delta \leq 1$ and $\beta < 1$.

We consider the following algorithm.

Input: A k -uniform hypergraph H , defined by the list of its edges, with a set V of km vertices.

Output: At least one of the following two (not mutually exclusive) conclusions:

- (a) The hypergraph H contains a matching with at least βm edges.
- (b) The hypergraph H contains at most $\delta^m \Phi_k(m)$ perfect matchings.

Algorithm: Let

$$\epsilon = \frac{1}{2} \delta^{1/(1-\beta)}.$$

Let us define a weight $W = \{w_S\}$ on the complete k -uniform hypergraph $\binom{V}{k}$ as follows:

$$(2.1.1) \quad w_S = \begin{cases} 1 & \text{if } S \in H \\ \epsilon & \text{if } S \notin H. \end{cases}$$

The weight W is ϵ^{-1} -balanced and we apply the algorithm of Theorem 1.5 to compute in polynomial in m time a number η such that

$$\eta \cdot m^{-\gamma} \leq P(W) \leq \eta \cdot m^\gamma$$

for some $\gamma = \gamma(\delta, \beta) > 0$.

If $m = 1$ or if

$$(2.1.2) \quad \frac{m}{\ln m} \leq \frac{2\gamma}{(1-\beta) \ln 2}$$

we check by direct enumeration whether (a) or (b) hold. Since k , β and δ are fixed in advance, this requires only a constant time.

If (2.1.2) does not hold, we output conclusion (a) if $\eta \cdot m^\gamma > \delta^m \Phi_k(m)$ and conclusion (b) if $\eta \cdot m^\gamma \leq \delta^m \Phi_k(m)$.

It is not hard to see that the algorithm is indeed correct. If $\eta \cdot m^\gamma \leq \delta^m \Phi_k(m)$ then $P(W) \leq \delta^m \Phi_k(m)$ and H necessarily contains not more than $\delta^m \Phi_k(m)$ perfect matchings. If $\eta \cdot m^\gamma > \delta^m \Phi_k(m)$ then, assuming that (2.1.2) does not hold, we conclude that

$$P(W) \geq \frac{\delta^m}{m^{2\gamma}} \Phi_k(m) > \frac{\delta^m}{2^{(1-\beta)m}} \Phi_k(m) = \epsilon^{(1-\beta)m} \Phi_k(m),$$

from which it follows that H contains a matching with not fewer than βm edges.

In particular, confronted with two hypergraphs on km vertices, one of which contains more than $\delta^m \Phi_k(m)$ perfect matchings and the other with no matchings of size βm or bigger, the algorithm will be able to decide which is which. It will necessarily output a) in the former case and b) in the latter.

(2.2) Definition. A k -uniform hypergraph H is called d -regular if every vertex of H is contained in exactly d edges of H .

For example, a complete k -uniform hypergraph with km vertices is d -regular for $d = \binom{km-1}{k-1}$. The existence of a perfect or nearly perfect matching in d -regular hypergraphs was extensively studied, see, for example, [Vu00] and references therein. As a corollary of Theorem 1.3, we obtain the following estimate for the number of nearly perfect matchings in a regular hypergraph.

(2.3) Theorem. *Let us fix an integer $k > 1$ and $0 < \alpha, \beta < 1$. Then there exists a positive integer $m_0 = m_0(k, \alpha, \beta)$ such that the following holds.*

Let H be a k -uniform d -regular hypergraph with km vertices where

$$d \geq \alpha \binom{km-1}{k-1} \quad \text{and} \quad m \geq m_0.$$

Then for every positive integer $s \leq \beta m$ the hypergraph H contains at least

$$\alpha^m \frac{\Phi_k(m)}{\Phi_k(m-s)}$$

matchings of size s .

Proof. All implied constants in the “ O ” notation below may depend on k, α and β only.

Let V be the set of vertices of a k -uniform d -regular hypergraph, $|V| = km$. Let us choose $0 < \epsilon < 1$ such that

$$(2.3.1) \quad \epsilon^{1-\beta} < \alpha + \epsilon(1 - \alpha)$$

and let us define a weight $W = \{w_S\}$ on the complete k -uniform hypergraph $\binom{V}{k}$ by (2.1.1). Then

$$\sum_{\substack{S \in \binom{V}{k} \\ S \ni v}} w_S = (1 - \epsilon)d + \epsilon \binom{km-1}{k-1} \quad \text{for all } v \in V.$$

It follows from Theorem 1.3 and scaling that

$$(2.3.2) \quad \begin{aligned} P(W) &\geq \left((1 - \epsilon)d + \epsilon \binom{km-1}{k-1} \right)^m e^{-m(k-1)} \frac{1}{m^{O(1)}} \\ &\geq (\alpha + \epsilon(1 - \alpha))^m \binom{km-1}{k-1}^m e^{-m(k-1)} \frac{1}{m^{O(1)}}. \end{aligned}$$

We note that

$$\begin{aligned}
(2.3.3) \quad \frac{\binom{km-1}{k-1}^m}{\Phi_k(m)} &= \frac{((km-1)!)^m (k!)^m m!}{((k-1)!)^m ((km-k)!)^m (km)!} = \frac{((km-1)!)^m k^m m!}{((km-k)!)^m (km)!} \\
&= \frac{((km)!)^m k^m m!}{(km)^m ((km-k)!)^m (km)!} = \frac{((km)!)^m m!}{((km-k)!)^m (km)! m^m} \\
&= \left(\frac{(km)(km-1) \cdots (km-k+1)}{(km)^k} \right)^m \cdot \frac{(km)^{km} m!}{(km)! m^m}.
\end{aligned}$$

Since $\ln(1-x) \geq -2x$ for $0 \leq x \leq 0.5$, we conclude that

$$\left(\frac{(km)(km-1) \cdots (km-k+1)}{(km)^k} \right)^m = \exp \left\{ m \sum_{i=1}^{k-1} \ln \left(1 - \frac{i}{km} \right) \right\} \geq e^{-k+1}.$$

Using Stirling's formula, we conclude from (2.3.3) and (2.3.2) that

$$(2.3.4) \quad P(W) \geq (\alpha + \epsilon(1-\alpha))^m \Phi_k(m) \frac{1}{m^{O(1)}}.$$

If a perfect matching in $\binom{V}{k}$ contains fewer than s edges of H then the contribution of the corresponding term to $P(W)$ is less than ϵ^{m-s} . Since every matching in H of size s can be appended to a perfect matching in $\binom{V}{k}$ in $\Phi_k(m-s)$ ways, we conclude that the number of matchings in H of size s is at least

$$\frac{P(W) - \epsilon^{m-s} \Phi_k(m)}{\Phi_k(m-s)} \geq \frac{P(W) - \epsilon^{(1-\beta)m} \Phi_k(m)}{\Phi_k(m-s)}.$$

The proof now follows from (2.3.4) and (2.3.1). \square

3. GENERAL RESULTS ON SCALING

In this section, we summarize some results on scaling which we need for the proofs of Theorems 1.3 and 1.5.

(3.1) Theorem. *Let H be a k -uniform hypergraph with a set V of $|V| = km$ vertices and let $\Omega_k(H)$ be the set of all k -stochastic weights on H . Suppose that the set $\Omega_k(H)$ has a non-empty relative interior, that is contains a positive weight $Y \in \Omega_k(H)$.*

For a positive weight $W = \{w_S : S \in H\}$ on H , let us define a function $f_W : \Omega_k(H) \rightarrow \mathbb{R}$ by

$$f_W(X) = \sum_{S \in H} x_S \ln \frac{x_S}{w_S} \quad \text{for } X \in \Omega_k(H), X = \{x_S : S \in H\}.$$

Then function f_W attains its minimum on $\Omega_k(H)$ at a unique weight $Z = \{z_S : S \in H\}$. We have $z_S > 0$ for all $S \in H$ and there exist real $\lambda_v > 0 : v \in V$ such that

$$(3.1.1) \quad z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all } S \in H.$$

We have

$$f_W(Z) = \sum_{v \in V} \ln \lambda_v.$$

Furthermore, if $\lambda_v > 0 : v \in V$ are reals such that weight Z defined by (3.1.1) is k -stochastic, then Z is the minimum point of f_W on $\Omega_k(H)$.

Proof. First, we observe that function f_W is strictly convex, so its minimum on the convex set $\Omega_k(H)$ is unique. Next,

$$(3.1.2) \quad \frac{\partial}{\partial x_S} f_W(X) = \ln \frac{x_S}{w_S} + 1,$$

which is finite if $x_S > 0$ and is $-\infty$ if $x_S = 0$ (we consider the right derivative in this case). If $z_S = 0$ for some S then for a sufficiently small $\epsilon > 0$ we have

$$f_W((1 - \epsilon)Z + \epsilon Y) < f_W(Z),$$

which is a contradiction. Hence $z_S > 0$ for all $S \in H$.

Since the minimum point Z lies in the relative interior of $\Omega_k(H)$, considered as a convex polyhedron in \mathbb{R}^H , the Lagrange multiplier condition implies that there exist real $\mu_v : v \in V$ such that

$$(3.1.3) \quad \ln \frac{z_S}{w_S} = \sum_{v \in S} \mu_v \quad \text{for all } S \in H.$$

Hence, letting $\lambda_v = e^{\mu_v}$ for $v \in V$, we obtain

$$z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all } S \in H.$$

Now,

$$f_W(Z) = \sum_{S \in H} z_S \left(\sum_{v \in S} \ln \lambda_v \right) = \sum_{v \in V} \ln \lambda_v \left(\sum_{\substack{S \in H \\ S \ni v}} z_S \right) = \sum_{v \in V} \ln \lambda_v,$$

as desired.

If (3.1.1) holds for some $\lambda_v > 0$ and k -stochastic $Z = \{z_S\}$, then (3.1.3) holds with $\mu_v = \ln \lambda_v$ and by (3.1.2) we conclude that Z is a critical point of f_W in the relative interior of $\Omega_k(H)$. Since f_W is strictly convex, Z must be the minimum point of f_W on $\Omega_k(H)$. \square

Theorem 3.1 implies that any positive weight W on a hypergraph H having a positive k -stochastic weight can be scaled uniquely to a k -stochastic weight Z , in which case we have $P_H(W) = \exp\{-f_W(Z)\}P_H(Z)$. Scaling factors $\{\lambda_v > 0 : v \in V\}$, however, do not have to be unique, as the example of a complete k -partite hypergraph readily shows (although in the case of the complete k -uniform hypergraph the scaling factors are unique). We note that if H is the complete k -uniform hypergraph or the complete k -partite hypergraph then there is a positive k -stochastic weight $Y = \{y_S : S \in H\}$ on H . In the former case we can choose

$$y_S = \binom{km - 1}{k - 1}^{-1} \quad \text{for all } S \in H,$$

while in the latter case we can choose

$$y_S = m^{-k+1} \quad \text{for all } S \in H.$$

We need a dual description of the scaling factors λ_v .

(3.2) Theorem. *Let H be a k -uniform hypergraph with a set V of $|V| = km$ vertices and let $W = \{w_S : S \in H\}$ be a positive weight on H . Let $\lambda_v > 0 : v \in V$ be reals such that the weight $Z = \{z_S\}$ defined by*

$$z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all } S \in H$$

is k -stochastic.

Let us define a set $C(W) \subset \mathbb{R}^V$ by

$$C(W) = \left\{ (x_v, v \in V) : \sum_{S \in H} w_S \exp \left\{ \sum_{v \in S} x_v \right\} \leq m \right\}.$$

Then the point $(\mu_v : v \in V)$, where $\mu_v = \ln \lambda_v$ for all $v \in V$, is a maximum point of the linear function $\sum_{v \in V} x_v$ on $C(W)$.

Proof. Since weight Z is k -stochastic, we have

$$\sum_{\substack{S \in H \\ S \ni u}} w_S \exp \left\{ \sum_{v \in S} \mu_v \right\} = 1 \quad \text{for all } u \in V,$$

which means that $(\mu_v : v \in V)$ is a critical point of the linear function $\sum_{v \in V} x_v$ on the smooth surface defined in \mathbb{R}^V by the equation

$$\sum_{S \in H} w_S \exp \left\{ \sum_{v \in S} x_v \right\} = m.$$

The set $C(W)$ is convex and hence $(\mu_v : v \in V)$ has to be an extremum point of function $\sum_{v \in V} x_v$ on $C(W)$. Moreover, it has to be a maximum point since the function is unbounded from below on $C(W)$. \square

4. SCALING BALANCED WEIGHTS

Our proof of Theorem 1.3 is based on two lemmas.

(4.1) Lemma. *Let us fix an integer $k > 1$ and a real $\alpha \geq 1$ and let H be a complete k -uniform hypergraph or a complete k -partite hypergraph. If $W = \{w_S\}$ is an α -balanced weight on H and if $Z = \{z_S\}$ is the k -stochastic weight obtained from W by scaling, then Z is α^{k+1} -balanced.*

Proof. Let V be the set of vertices of hypergraph H . Without loss of generality, we assume that $|V| > k$. For a subset $X \subset V$, we denote by

$$H_X = \{S \in H : S \supset X\}$$

the set of edges of H containing X . Let $\{\lambda_v > 0 : v \in V\}$ be scaling factors so that

$$(4.1.1) \quad z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all } S \in H.$$

Suppose first that $H = V_1 \times \dots \times V_k$ is a complete k -partite hypergraph, so $V = V_1 \cup \dots \cup V_k$ and $|V_1| = \dots = |V_k|$. For every $i = 1, \dots, k$ and for every pair of vertices $v, u \in V_i$ we have

$$(4.1.2) \quad \sum_{S \in H_{\{v\}}} z_S = \sum_{S \in H_{\{u\}}} z_S = 1.$$

Let us consider the bijection $\phi : H_{\{v\}} \rightarrow H_{\{u\}}$ defined by

$$(4.1.3) \quad \phi(S) = S \cup \{u\} \setminus \{v\}.$$

By (4.1.1) we have

$$(4.1.4) \quad \frac{z_{\phi(S)}}{z_S} = \frac{\lambda_u}{\lambda_v} \cdot \frac{w_{\phi(S)}}{w_S}.$$

Since weight W is α -balanced, in view of (4.1.2) we conclude that

$$(4.1.5) \quad \frac{1}{\alpha} \leq \frac{\lambda_u}{\lambda_v} \leq \alpha,$$

which proves that Z is α^{k+1} -balanced.

Suppose now that $H = \binom{V}{k}$ is a complete k -uniform hypergraph. Then for any two distinct vertices $u, v \in V$ we have

$$(4.1.6) \quad \sum_{S \in H_v \setminus H_{\{u,v\}}} z_S = \sum_{S \in H_u \setminus H_{\{u,v\}}} z_S = 1 - \sum_{S \in H_{\{u,v\}}} z_S > 0.$$

Let us consider the bijection $\phi : H_v \setminus H_{\{u,v\}} \rightarrow H_u \setminus H_{\{u,v\}}$ defined by (4.1.3). From (4.1.1) we deduce that (4.1.4) holds and in view of (4.1.6) we conclude that (4.1.5) follows. Since weight W is α -balanced, (4.1.5) implies that Z is α^{k+1} -balanced. \square

The second lemma asserts that if we scale to a k -stochastic weight a weight which is already sufficiently close to being k -stochastic, then the product of the scaling factors is close to 1.

(4.2) Lemma. *Let us fix an integer $k > 1$ and a real $\alpha \geq 1$. Then there exist a positive integer $m_0 = m_0(k, \alpha)$ and a real $\beta = \beta(k, \alpha) > 0$ such that the following holds.*

Let H be the complete k -uniform hypergraph or a complete k -partite hypergraph with a set V of vertices, where $|V| = km$ for some integer $m \geq m_0$. Suppose that $W = \{w_S\}$ is an α -balanced weight on H , that

$$\sum_{S \in H} w_S = m$$

and that

$$\sum_{\substack{S \in H \\ S \ni v}} w_S = 1 - \delta_v \quad \text{where} \quad |\delta_v| \leq \frac{\alpha}{m} \quad \text{for all} \quad v \in V.$$

Let $\lambda_v > 0 : v \in V$ be reals such that weight $Z = \{z_S\}$ defined by

$$z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all} \quad S \in H$$

is k -stochastic. Then

$$0 \leq \sum_{v \in V} \ln \lambda_v \leq \frac{\beta}{m}.$$

Proof. All implied constants in the “ O ” notation below may depend only on k and α .

We note that the point $(x_v = 0 : v \in V)$ belongs to the set $C(W)$ of Theorem 3.2, and so by Theorem 3.2 we have

$$\sum_{v \in V} \ln \lambda_v \geq \sum_{v \in V} x_v = 0.$$

Next, we note that

$$(4.2.1) \quad \sum_{v \in V} \delta_v = \sum_{v \in V} \left(1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \right) = km - k \sum_{S \in H} w_S = 0.$$

In addition, if $H = V_1 \times \dots \times V_k$ is the complete k -partite graph, where $V = V_1 \cup \dots \cup V_k$ and $|V_1| = \dots = |V_k| = m$, for every $i = 1, \dots, k$ we have

$$(4.2.2) \quad \sum_{v \in V_i} \delta_v = \sum_{v \in V_i} \left(1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \right) = m - \sum_{S \in H} w_S = 0.$$

We define numbers $\{\rho_S : S \in H\}$ as follows. If $H = \binom{V}{k}$ is the complete k -uniform hypergraph, we define

$$\rho_S = \binom{km-2}{k-1}^{-1} \sum_{v \in S} \delta_v \quad \text{for all } S \in H.$$

If $H = V_1 \times \dots \times V_k$ is a complete k -partite graph, we define

$$\rho_S = \frac{1}{m^{k-1}} \sum_{v \in S} \delta_v \quad \text{for all } S \in H.$$

We claim that

$$(4.2.3) \quad \sum_{\substack{S \in H \\ S \ni v}} \rho_S = \delta_v \quad \text{for all } v \in V.$$

Indeed, if $H = \binom{V}{k}$ then using (4.2.1) we obtain

$$\begin{aligned} \sum_{\substack{S \in H \\ S \ni v}} \rho_S &= \frac{\binom{km-1}{k-1}}{\binom{km-2}{k-1}} \delta_v + \frac{\binom{km-2}{k-2}}{\binom{km-2}{k-1}} \sum_{u \in V \setminus \{v\}} \delta_u \\ &= \frac{\binom{km-1}{k-1} - \binom{km-2}{k-2}}{\binom{km-2}{k-1}} \delta_v \\ &= \delta_v \end{aligned}$$

and if $H = V_1 \times \dots \times V_k$ then using (4.2.2) we obtain that for all $i = 1, \dots, k$ and for all $v \in V_i$ we have

$$\sum_{\substack{S \in H \\ S \ni v}} \rho_S = \delta_v + \frac{m^{k-2}}{m^{k-1}} \sum_{u \in V \setminus V_i} \delta_u = \delta_v.$$

In either case, (4.2.3) holds. In addition, from (4.2.1)

$$(4.2.4) \quad \sum_{S \in H} \rho_S = \frac{1}{k} \sum_{v \in V} \sum_{\substack{S \in H \\ S \ni v}} \rho_S = \frac{1}{k} \sum_{v \in V} \delta_v = 0.$$

Let us define

$$x_S = w_S + \rho_S \quad \text{for all } S \in H.$$

Then, from (4.2.3) we have

$$(4.2.5) \quad \sum_{\substack{S \in H \\ S \ni v}} x_S = 1 \quad \text{for all } v \in V.$$

Since weight W is α -balanced, we have

$$w_S \geq \frac{m}{\alpha |H|} \quad \text{for all } S \in H.$$

In particular,

$$(4.2.6) \quad \min_{S \in H} w_S \geq \gamma m^{k-1} \quad \text{for some } \gamma = \gamma(k, \alpha) > 0.$$

On the other hand, for all $S \in H$

$$|\rho_S| \leq \binom{km-2}{k-1}^{-1} \frac{\alpha k}{m} \quad \text{when } H = \binom{V}{k}$$

and

$$|\rho_S| \leq \frac{\alpha k}{m^k} \quad \text{when } H = V_1 \times \dots \times V_k.$$

In either case,

$$(4.2.7) \quad \max_{S \in H} |\rho_S| = O(m^{-k}).$$

From (4.2.6) and (4.2.7) we conclude that

$$x_S \geq 0 \quad \text{for all } S \in H \quad \text{provided } m \geq m_0(k, \alpha).$$

Therefore, from (4.2.5) we conclude $X = \{x_S\}$ is a k -stochastic weight on H provided $m \geq m_0(k, \alpha)$. Using (4.2.4), (4.2.6) and (4.2.7), we conclude from Theorem 3.1 that for $m \geq m_0(k, \alpha)$ we have

$$\begin{aligned} \sum_{v \in V} \ln \lambda_v &\leq \sum_{S \in H} x_S \ln \frac{x_S}{w_S} = \sum_{S \in H} (w_S + \rho_S) \ln \frac{w_S + \rho_S}{w_S} \\ &= \sum_{S \in H} (w_S + \rho_S) \ln \left(1 + \frac{\rho_S}{w_S} \right) \leq \sum_{S \in H} (w_S + \rho_S) \frac{\rho_S}{w_S} \\ &= \sum_{S \in H} \frac{\rho_S^2}{w_S} = O\left(\frac{1}{m}\right), \end{aligned}$$

which completes the proof. \square

5. PROOFS OF THEOREMS 1.3 AND 1.5

Our approach is somewhat similar to Bregman's original approach [Br73] combining scaling and induction to obtain upper bounds on permanents.

(5.1) Proof of Theorem 1.3. All implied constants in the “ O ” notation below may depend on k and α only.

Let H be either a complete k -uniform hypergraph $\binom{V}{k}$ or a complete k -partite graph $V_1 \times \dots \times V_k$ with a set V of $|V| = km$ vertices. Let $Z = \{z_S\}$ be a k -stochastic α -balanced weight on H . If $H = \binom{V}{k}$ and $U \subset V$ is a subset such that $|U| = kl$ for some integer $l \geq 1$, we consider the induced hypergraph $H|U$ consisting of the edges $S \in H$ such that $S \subset U$. Hence $H|U = \binom{U}{k}$ is the complete k -uniform hypergraph with the set U of vertices. Similarly, if $H = V_1 \times \dots \times V_k$ and if $|U \cap V_1| = \dots = |U \cap V_k| = l$ for some integer $l \geq 1$, we consider the restriction $H|U$ consisting of the edges $S \in H$ such that $S \subset U$. In this case, $H|U$ is a uniform k -partite graph with the set U of vertices, $H|U = U_1 \times \dots \times U_k$ where $U_i = V_i \cap U$ for $i = 1, \dots, k$.

For a subset $U \subset V$ as above, we define a weight

$$Z^U = \{z_S^U : S \in H|U\}$$

on $H|U$ as follows. We consider the restriction of weight Z onto hypergraph $H|U$ and define Z^U to be the scaling of the restriction to a k -stochastic weight. Since the restriction of Z onto $H|U$ is α -balanced, by Lemma 4.1 the weight Z^U is α^{k+1} -balanced. We consider the partition function associated with the hypergraph $H|U$, which we denote by P_U .

Our goal is to prove that

$$(5.1.1) \quad P_U(Z^U) = e^{-(k-1)l} \exp \left\{ O \left(\sum_{j=1}^l \frac{1}{j} \right) \right\} \quad \text{where } |U| = kl.$$

Applied to $l = m$, $U = V$ and $Z^U = Z$, formula (5.1.1) gives the desired result.

We prove (5.1.1) by induction on l .

Let us pick an element $u \in U$. Then there is a recursion

$$(5.1.2) \quad P_U(Z^U) = \sum_{\substack{A \in H|U \\ A \ni u}} z_A^U \cdot P_{U \setminus A}(Z^U).$$

Here $P_{U \setminus A}(Z^U)$ is the partition function computed on the restriction of the weight Z^U onto the hypergraph $H|(U \setminus A)$, which is not the same as $P_{U \setminus A}(Z^{U \setminus A})$, since the weight $Z^{U \setminus A}$ is obtained from Z^U by restricting it onto $H|(U \setminus A)$ and scaling the restriction to a k -stochastic weight.

Since Z^U is a k -stochastic weight on $H|U$, we have

$$(5.1.3) \quad \sum_{\substack{A \in H|U \\ A \ni u}} z_A^U = 1 \quad \text{and} \quad z_A^U \geq 0 \quad \text{for all} \quad A \in H|U.$$

Let

$$\sigma_A^U = \sum_{S \in H|(U \setminus A)} z_S^U$$

be the sum of the weights in the restriction Z^U onto $H|(U \setminus A)$, that is, the sum of the weights z_S^U for the edges $S \subset U$ not intersecting an edge $A \in H|U$. Since weight Z^U is k -stochastic and α^{k+1} -balanced, we have

$$(5.1.4) \quad \sigma_A^U = l - k + O\left(\frac{1}{l}\right) \quad \text{where} \quad |U| = kl.$$

Let us define a weight

$$W^{U \setminus A} = \left\{ w_S^{U \setminus A} : S \in H|(U \setminus A) \right\}$$

by scaling the restriction of the weight Z^U onto $H|(U \setminus A)$ to the total sum $l - 1$, so that

$$(5.1.5) \quad w_S^{U \setminus A} = \frac{l - 1}{\sigma_A^U} z_S^U \quad \text{for all} \quad S \in H|(U \setminus A).$$

Hence

$$(5.1.6) \quad P_{U \setminus A}(Z^U) = \left(\frac{\sigma_A^U}{l - 1} \right)^{l-1} P_{U \setminus A}(W^{U \setminus A}).$$

From (5.1.4), we have

$$(5.1.7) \quad \left(\frac{\sigma_A^U}{l - 1} \right)^{l-1} = \exp \left\{ -k + 1 + O\left(\frac{1}{l}\right) \right\}.$$

Since Z^U is k -stochastic and α^{k+1} -balanced, from (5.1.4) and (5.1.5) we conclude that

$$(5.1.8) \quad \max_{v \in U \setminus A} \left| \sum_{\substack{S \in H(U \setminus A) \\ S \ni v}} w_S^{U \setminus A} - 1 \right| = O\left(\frac{1}{l}\right).$$

We intend to apply Lemma 4.2 to weight $W^{U \setminus A}$. We observe that weight $W^{U \setminus A}$ is obtained from weight Z^U by restricting it onto the set $U \setminus A$ and then scaling by (5.1.5) to the total sum $l - 1$ of components. Therefore, the k -stochastic weight on $U \setminus A$ obtained from $W^{U \setminus A}$ by scaling is just $Z^{U \setminus A}$, the k -stochastic weight obtained by restricting the original weight Z onto $U \setminus A$ and scaling. Since $W^{U \setminus A}$ is α^{k+1} -balanced and (5.1.8) holds, by Lemma 4.2 we conclude that

$$(5.1.9) \quad P_{U \setminus A}(W^{U \setminus A}) = \exp\left\{O\left(\frac{1}{l}\right)\right\} P_{U \setminus A}(Z^{U \setminus A})$$

provided $l \geq m_0$ for some integer $m_0 = m_0(k, \alpha)$.

Combining (5.1.2), (5.1.6), (5.1.7) and (5.1.9), we conclude that

$$(5.1.10) \quad P_U(Z^U) = \sum_{\substack{A \in H|U \\ A \ni u}} z_A^U \exp\left\{-k + 1 + O\left(\frac{1}{l}\right)\right\} P_{U \setminus A}(Z^{U \setminus A})$$

provided $l \geq m_0(k, \alpha)$. The proof of (5.1.1) now follows from (5.1.10) and (5.1.3) by induction on l . \square

(5.2) Proof of Theorem 1.5. Let Z be the k -stochastic weight obtained from weight W by scaling. By Theorem 3.1 we have

$$P_H(Z) = f_W(Z)P_H(W) = e^\zeta P_H(W).$$

Moreover, by Lemma 4.1, weight Z is α^{k+1} -balanced and the proof follows by Theorem 1.3 applied to Z . \square

REFERENCES

- [AF08] N. Alon and S. Friedland, *The maximum number of perfect matchings in graphs with a given degree sequence*, Note 13, 2 pp., Electron. J. Combin. **15** (2008).
- [Br73] L.M. Bregman, *Certain properties of nonnegative matrices and their permanents. (Russian)*, Dokl. Akad. Nauk SSSR **211** (1973), 27–30.
- [D87a] S.J. Dow and P.M. Gibson, *An upper bound for the permanent of a 3-dimensional (0, 1)-matrix*, Proc. Amer. Math. Soc. **99** (1987), 29–34.
- [D87b] S.J. Dow and P.M. Gibson, *Permanents of d -dimensional matrices*, Linear Algebra Appl. **90** (1987), 133–145.

- [Eg81] G.P. Egorychev, *The solution of van der Waerden's problem for permanents*, Adv. in Math. **42** (1981), 299–305.
- [Fa81] D.I. Falikman, *Proof of the van der Waerden conjecture on the permanent of a doubly stochastic matrix. (Russian)*, Mat. Zametki **29** (1981), 931–938.
- [Fr10] S. Friedland, *Positive diagonal scaling of a nonnegative tensor to one with prescribed slice sums*, preprint [arXiv:0908.2368](https://arxiv.org/abs/0908.2368) (2010).
- [Gu08] L. Gurvits, *Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all. With a corrigendum*, Research Paper 66, 26 pp., Electron. J. Combin. **15** (2008).
- [J+04] M. Jerrum, A. Sinclair and E. Vigoda, *A polynomial-time approximation algorithm for the permanent of a matrix with nonnegative entries*, J. ACM **51** (2004), 671–697.
- [Ka72] R.M. Karp, *Reducibility among combinatorial problems*, Complexity of Computer Computations (Proc. Sympos., IBM Thomas J. Watson Res. Center, Yorktown Heights, N.Y., 1972), Plenum, New York, 1972, pp. 85–103.
- [L+00] N. Linial, A. Samorodnitsky and A. Wigderson, *A deterministic strongly polynomial algorithm for matrix scaling and approximate permanents*, Combinatorica **20** (2000), 545–568.
- [LP09] L. Lovász and M.D. Plummer, *Matching Theory*, AMS Chelsea Publishing, Providence, RI, 2009.
- [Mi78] H. Minc, *Permanents*, Encyclopedia of Mathematics and its Applications, Vol. 6, Addison-Wesley Publishing Co., Reading, Mass., 1978.
- [NN94] Y. Nesterov and A. Nemirovskii, *Interior-Point Polynomial Algorithms in Convex Programming*, SIAM Studies in Applied Mathematics, 13, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1994.
- [Sc78] A. Schrijver, *A short proof of Minc's conjecture*, J. Combinatorial Theory Ser. A **25** (1978), 80–83.
- [So03] G.W. Soules, *New permanental upper bounds for nonnegative matrices*, Linear Multilinear Algebra **51** (2003), 319–337.
- [Va79] L.G. Valiant, *The complexity of computing the permanent*, Theoret. Comput. Sci. **8** (1979), 189–201.
- [Vu00] V.H. Vu, *New bounds on nearly perfect matchings in hypergraphs: higher codegrees do help*, Random Structures Algorithms **17** (2000), 29–63.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043, USA

E-mail address: `barvinok@umich.edu`

DEPARTMENT OF COMPUTER SCIENCE, HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM CAMPUS, 91904, ISRAEL

E-mail address: `salex@cs.huji.ac.il`