

Multi-linear Systems and Invariant Theory
in the Context of Computer Vision and Graphics

Class 5: Self Calibration

CS329
Stanford University

Amnon Shashua

Material We Will Cover Today

- The basic equations and counting arguments
- The “absolute conic” and its image.
- Kruppa’s equations
- Recovering internal parameters.

The Basic Equations and Counting Arguments

Recall,

$$p \cong K[R;T] \begin{pmatrix} U \\ V \\ W \\ 1 \end{pmatrix} \quad \begin{matrix} \text{3D} \rightarrow \text{2D from Euclidean world frame to image} \\ \\ \\ \end{matrix}$$

$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = R \begin{pmatrix} U \\ V \\ W \end{pmatrix} + T \quad \text{world frame to first camera frame}$$

Let K, K' be the internal parameters of camera 1,2 and choose canonical frame in which $R=I$ and $T=0$ for first camera.

$$p \cong [K; \quad 0]P$$

$$p' \cong K' [R \quad t]P$$

$$P = \begin{pmatrix} X \\ Y \\ Z \\ 1 \end{pmatrix}$$

The Basic Equations and Counting Arguments

$$p \cong [I; \quad 0] W W^{-1} \begin{pmatrix} p \\ \mu \end{pmatrix} \qquad p = \frac{1}{Z} K \begin{pmatrix} X \\ Y \\ Z \end{pmatrix}$$

$$\text{where } \begin{pmatrix} X \\ Y \\ Z \\ 1 \end{pmatrix} \cong W^{-1} \begin{pmatrix} p \\ \mu \end{pmatrix}$$

$$W = \begin{bmatrix} K & 0 \\ n^T & 1 \end{bmatrix}$$

(8 unknown parameters)

W^{-1} maps from the projective frame to Euclidean

The Basic Equations and Counting Arguments

$(X, Y, Z, 0)$ are the points on the plane at infinity (in Euc frame)

→ $(0,0,0,1)$ is the plane at infinity


$$\pi_{\infty} = W^{-T} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \text{ is the plane at infinity in Proj frame}$$

(recall: if W maps points to points (Euc \rightarrow Proj), then the dual

W^{-T} maps planes to planes)

The Basic Equations and Counting Arguments

$$\pi_\infty = W^{-T} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{bmatrix} K^{-T} & -K^{-T}n \\ 0^T & 1 \end{bmatrix} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -K^{-T}n \\ 1 \end{pmatrix} = \begin{pmatrix} u \\ 1 \end{pmatrix}$$

 $n = -K^T u$

The Basic Equations and Counting Arguments

$$p' \cong [H \quad e'] \begin{pmatrix} p \\ \mu \end{pmatrix} \quad \text{Projective frame}$$

$$= [H \quad e'] W W^{-1} \begin{pmatrix} p \\ \mu \end{pmatrix}$$

$$= [H \quad e'] \begin{bmatrix} K & 0 \\ n^T & 1 \end{bmatrix} W^{-1} \begin{pmatrix} p \\ \mu \end{pmatrix}$$

$$= K' [R \quad t] W^{-1} \begin{pmatrix} p \\ \mu \end{pmatrix}$$

$$\longrightarrow K' [R \quad t] \cong [H \quad e'] \begin{bmatrix} K & 0 \\ n^T & 1 \end{bmatrix}$$

The Basic Equations and Counting Arguments

$$K' [R \quad t] \cong [H \quad e'] \begin{bmatrix} K & 0 \\ n^T & 1 \end{bmatrix}$$

→ $HK + e' n^T \cong K' R$

$$K'^{-1} HK + K'^{-1} e' n^T \cong R$$

since $n = -K^T u$ then,

$$K'^{-1} HK - K'^{-1} e' u^T K \cong R$$

Class 5 but $RR^T \cong I$ provides 5 (non-linear) constraints!

The Basic Equations and Counting Arguments

$$K^{-1}HK - K^{-1}e'u^TK \cong R$$

$$RR^T \cong I \quad \rightarrow$$

$$(H - e'u^T)KK^T(H - e'u^T)^T \cong K'K'^T$$

Since the right-hand side is symmetric and up to scale, we have 5 constraints.

The Basic Equations and Counting Arguments

Lets do some counting:

Let $1 \leq \# K \leq 5$ be the number of internal parameters

m be the number of views

$$5(m - 1) \geq (\# K) + (m - 1)(\# K) + 3$$

The Basic Equations and Counting Arguments

$$5(m - 1) \geq (\# K) + (m - 1)(\# K) + 3$$

$\# K = 5 \longrightarrow$ not enough measurements (!)

$\# K = 4 \longrightarrow 5(m - 1) \geq 7 + 4(m - 1) \longrightarrow m \geq 8$

$\# K = 3 \longrightarrow 5(m - 1) \geq 6 + 3(m - 1) \longrightarrow m \geq 4$

$\# K = 1 \longrightarrow 5(m - 1) \geq 4 + (m - 1) \longrightarrow m \geq 2$

$K = K' = K'' \dots \longrightarrow 5(m - 1) \geq 8 \longrightarrow m \geq 3$
(fixed internal params)

The remainder of this lecture is about a geometric insight of

$$(H - e'e^T)KK^T(H - e'e^T)^T \cong K'K'^T$$


The Absolute Conic

$p^T C p = 0$ where $C = C^T$ represents a conic in 2D

$(x_1, x_2, x_3, 0)$ are the points on the plane at infinity (in Euc frame)

 $(0,0,0,1)$ is the plane at infinity

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}^T C \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \text{is conic on the plane at infinity}$$

when $C = I$  $x_1^2 + x_2^2 + x_3^2 = 0$

Class 5 is the “absolute” conic (imaginary circle) Ω_∞

The Absolute Conic

Plane at infinity is preserved under affine transformations:

$$W = \begin{bmatrix} A & v \\ 0^T & 1 \end{bmatrix} \quad \text{because} \quad W \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ 0 \end{pmatrix} = \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \\ 0 \end{pmatrix}$$

Ω_∞ is preserved under similarity transformation (R,t up to scale)

$$W \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ 0 \end{pmatrix} = A \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \quad \text{if} \quad p^T C p = 0 \quad \text{and} \quad A p \cong p'$$

$$\text{then } p'^T C' p' = p^T (A^T C' A) p = 0$$

$$\longrightarrow C' = A^{-T} C A^{-1}$$

but $C = I$, so in order that $C' = I$ we must have:

$$\text{Class 5} \quad I \cong A^{-T} A^{-1} \longrightarrow A \text{ is orthogonal}$$

The Image of the Absolute Conic

Image of points at infinity:

$$K [R \quad t] \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ 0 \end{pmatrix} = KR \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

let $A = KR$

if $p^T C p = 0$ is a conic on the plane at infinity

then $C' = A^{-T} C A^{-1}$ is the projected conic onto the image

since $C = I$, then $C' = (KR)^{-T} (KR)^{-1} = K^{-T} K^{-1}$

→ the image of Ω_∞ is

$$\omega = K^{-T} K^{-1}$$

The Image of the Dual Absolute Conic

$p^T C p = 0 \quad \longrightarrow \quad l = C p \quad \text{is tangent to the conic at } p$

$$p = C^{-1} l \quad \longrightarrow \quad p^T C p = l^T C^{-T} C C^{-1} l = l^T C^{-1} l = 0$$

$\longrightarrow \quad \boxed{\omega^* = K K^T}$ is the image of the dual absolute conic

The basic equation:


$$(H - e' u^T) K K^T (H - e' u^T)^T \cong K' K'^T$$

Becomes:

$$(H - e' u^T) \omega^* (H - e' u^T)^T \cong \omega'^*$$

Why 8 parameters? 5 for the conic, 3 for the plane

Geometric Interpretation of ω



$$\begin{pmatrix} d \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

$$p \cong \begin{bmatrix} K & 0 \end{bmatrix} \begin{pmatrix} \lambda d \\ 1 \end{pmatrix} \cong Kd \quad \longrightarrow \quad d = K^{-1} p \quad \text{direction of optical ray}$$

The angle between two optical rays d_1, d_2

$$\cos(\theta) = \frac{d_1^T d_2}{|d_1| |d_2|} = \frac{p_1^T \omega p_2}{\sqrt{p_1^T \omega p_1} \sqrt{p_2^T \omega p_2}}$$

\longrightarrow given $\omega = K^{-T} K^{-1}$ one can measure angles


Kruppa's Equations

General idea: eliminate n from the basic equation.

$$p_i^T C p_i = 0$$

$$l_i^T C^{-1} l_i = 0$$

$$l_i = [e]_{\times} p_i$$



$$p_i^T [e]_{\times} C^{-1} [e]_{\times} p_i = p_i^T C_t p_i = 0$$

$$p_i'^T [e']_{\times} C'^{-1} [e']_{\times} p_i' = p_i'^T C_t' p_i' = 0$$

Class 5 C_t, C_t' are degenerate (rank 2) conics

Kruppa's Equations

Note:

$$C_t = l_1 l_2^T + l_2 l_1^T \quad \text{is a degenerate conic}$$

$$p^T C_t p = 0 \quad \text{iff} \quad p^T l_1 = 0 \quad \text{or} \quad p^T l_2 = 0$$

Let H_π be the homography induced by the plane of the conic

$$C_t \cong H^T C'_t H \quad (\text{slide 14})$$

$$\begin{array}{c} \rightarrow \\ [e]_\times C^{-1} [e]_\times \cong H_\pi^T [e']_\times C'^{-1} [e']_\times H_\pi \\ \underbrace{\hspace{10em}}_{C_t} \qquad \underbrace{\hspace{10em}}_{C'_t} \end{array}$$

Kruppa's Equations

$$[e]_{\times} C^{-1} [e]_{\times} \cong H_{\pi}^T [e']_{\times} C'^{-1} [e']_{\times} H_{\pi}$$

Recall: $F = [e']_{\times} H_{\pi}$

$\rightarrow [e]_{\times} C^{-1} [e]_{\times} \cong F^T C'^{-1} F$

In our case $\pi = \infty$ and the conic is Ω_{∞}

$\rightarrow [e]_{\times} \omega^* [e]_{\times} \cong F^T \omega'^* F$

$[e]_{\times} K K^T [e]_{\times} \cong F^T K' K'^T F$

Likewise:

$[e']_{\times} \omega'^* [e']_{\times} \cong F \omega^* F^T$

Determining K given π_∞

Recall: $\pi_\infty = \begin{pmatrix} u \\ 1 \end{pmatrix}$ the location of the plane at infinity in the projective coordinate frame.

We wish to represent the homography H_∞ induced by π_∞

Let $P = \begin{pmatrix} p \\ \mu \end{pmatrix}$ $P^T \pi_\infty = 0$ be a point on the plane at infinity.

$$\rightarrow p^T u + \mu = 0 \quad \rightarrow P = \begin{pmatrix} p \\ -p^T u \end{pmatrix}$$

$$p' \cong [H \quad e'] \begin{pmatrix} p \\ \mu \end{pmatrix} = Hp - (p^T u)e' = (H - e'u^T)p$$

Class 5



$$H_\infty \cong H - e'u^T$$

Determining K given π_∞

Recall: $(H - e'u^T)\omega^*(H - e'u^T)^T \cong \omega'^*$ (slide 16)

$$\omega^* = KK^T \quad \omega'^* = K'K'^T$$



$$H_\infty \omega^* H_\infty^T \cong \omega'^*$$

Note: this could be derived from “first principles” as well:

$$l^T \omega^* l = 0$$

$$l'^T \omega'^* l' = 0$$

tangents lines to the image of the absolute conic

$$H_\infty^{-T} l \cong l' \quad \longrightarrow \quad l^T \underbrace{H_\infty^{-1} \omega'^* H_\infty^{-T}}_{\omega^*} l = 0$$

Determining K given π_∞

$$H_\infty \omega^* H_\infty^T \cong \omega'^*$$

Assume fixed internal parameters $\omega^* = \omega'^*$

$$H_\infty \omega^* H_\infty^T = \omega^* \quad \det(H_\infty) = 1$$


Note: $\det(H_\infty) = 1 \rightarrow H_\infty = KRK^{-1}$

$$KRK^{-1}KK^TK^{-T}R^TK^T = KK^T$$

Provides 4 independent linear constraints on $\omega^* = KK^T$

Why 4 and not 5?

Class 5

 we need 3 views (since ω^* has 5 unknowns²³)

Why 4 Constraints?

$$H_{\infty} \omega^* H_{\infty}^T = \omega^*$$

$$\det(H_{\infty}) = 1 \quad \longrightarrow \quad H_{\infty} = K R K^{-1}$$

H_{∞} and R are “similar” matrices, i.e., have the same eigenvalues

Let w be the axis of rotation, i.e., $Rw = w$

\longrightarrow H_{∞} has an eigenvalue = 1, with eigenvector $v = Kw$

$$(K R K^{-1} v = v \quad \longrightarrow \quad R(K^{-1} v) = K^{-1} v)$$

Why 4 Constraints?

$$H_{\infty} v = v$$

if ω^* is a solution to $H_{\infty} \omega^* H_{\infty}^T = \omega^*$

then $\omega^* + \lambda v v^T$ is also a solution

$$\begin{aligned} H_{\infty} (\omega^* + \lambda v v^T) H_{\infty}^T &= H_{\infty} \omega^* H_{\infty}^T + \lambda (H_{\infty} v)(H_{\infty} v)^T \\ &= \omega^* + \lambda v v^T \end{aligned}$$



We need one more camera motion (with a different axis of rotation).

Kruppa's Equations (revisited)

Kruppa's equations: $[e']_{\times} \omega'^* [e']_{\times} \cong F \omega^* F^T$

Start with the basic equation:

$$H_{\infty} \omega^* H_{\infty}^T \cong \omega'^*$$

Multiply the terms by $[e']_{\times}$ on both sides:

$$[e']_{\times} \omega'^* [e']_{\times} \cong \underbrace{[e']_{\times} H_{\infty} \omega^* H_{\infty}^T [e']_{\times}}_F$$