

## GIRTH AND EUCLIDEAN DISTORTION

N. LINIAL, A. MAGEN AND A. NAOR

### Abstract

Let  $G$  be a  $k$ -regular graph,  $k \geq 3$ , with girth  $g$ . We prove that every embedding  $f : G \rightarrow \ell_2$  has distortion  $\Omega(\sqrt{g})$ . Two proofs are given, one based on Markov type [B] and the other on quadratic programming. In the core of both proofs are some Poincaré-type inequalities on graph metrics.

### 1 Introduction

Finite metric spaces and their embeddings in other metric spaces have been intensively investigated in recent years. For metric spaces  $(X, d_X)$  and  $(Y, d_Y)$ , and an embedding  $f : X \rightarrow Y$  we define the distortion of  $f$  by

$$\text{dist}(f) = \sup_{x,y \in X} \frac{d_Y(f(x), f(y))}{d_X(x, y)} \cdot \sup_{x,y \in X} \frac{d_X(x, y)}{d_Y(f(x), f(y))}.$$

We denote by  $c_Y(X, d)$  the least distortion with which  $(X, d)$  may be embedded in  $Y$ . For  $p \geq 1$  we denote  $c_p(X, d) = c_{\ell_p}(X, d)$ . A special case of interest is when  $Y$  is the Euclidean space  $\ell_2$ . In this case, a fundamental result of Bourgain [Bou1] states that  $c_2(X, d) = O(\log n)$  for every  $n$ -point metric space  $(X, d)$ .

One natural source for examples of metrics comes from graphs. A graph  $G$  induces a metric  $d_G$  on its vertex set, where  $d_G(u, v)$  is the length of the shortest path in  $G$  joining  $u$  and  $v$ . Special families of graphs define special families of metrics, e.g. expander graph metrics are studied in [LLR], [M1], [LM], tree metrics in [Ba], [Bou2], [LS], [M2], metrics of graphs with forbidden minors in [R], [GNRS], [KPR] and many more. Here we consider regular graphs with constant degree, and wish to study the Euclidean distortion of these graphs as a function of their girth.

---

N.L. was supported in part by grants from the Israel Science Foundation and the Binational Science Foundation Israel-USA. A.N. was supported in part by the Binational Science Foundation Israel-USA and the Clore Foundation. This work is part of a Ph.D. thesis being prepared under the supervision of Professor Joram Lindenstrauss.

In [LLR], Bourgain's upper bound for the Euclidean distortion was shown to be tight. In fact, the Euclidean distortion of an  $n$  point constant degree expander is  $\Omega(\log n)$ . This fact is striking since for any graph  $G$ , the trivial bound  $c_2(G, d_G) \leq \text{diam}(G)$  follows by embedding  $G$  as a simplex in  $\ell_2$ . When  $G$  is an  $n$ -point expander,  $\text{diam}(G) = O(\log n)$ , so that up to constant factors, the best way to embed an expander in Hilbert space is to ignore its structure altogether and embed it as if it were a clique! A new proof of this phenomenon follows from the results presented in this article.

In most examples we know, metrics are far from being Euclidean, since they include "too many" triples for which the triangle inequality holds as (a near) equality. The simplest example is  $K_{1,3}$  that cannot be embedded isometrically in Euclidean space, since there cannot be three geodesics between three different points that meet in their interior. For the hypercube of dimension  $m$ , there are  $m!$  geodesics between every pair of antipodes, and consequently we get a large Euclidean distortion [E], [LM].

It is known that for a tree  $T$  on  $n$  vertices,  $c_2(T, d_T) = O(\sqrt{\log \log n})$ , and this bound is tight, as the Euclidean distortion of the complete binary tree on  $n$  vertices is  $\Theta(\sqrt{\log \log n})$  (see [Bou2], [M2], [LS]). Motivated by this, Linial, London and Rabinovich considered graphs with vertex degrees bigger than 2 and with girth  $g$ . Any such graph contains isometrically a tree of depth  $g/2 - 1$ , which immediately gives the lower bound  $c_2(G, d_G) = \Omega(\sqrt{\log g})$ . Unlike the complete binary tree example, every vertex in the graph is a root of such a tree. In [LLR], Linial et al. conjecture that  $c_2(d_G) = \Omega(g)$ . In this paper we prove  $c_2(d_G) = \Omega(\sqrt{g})$  for  $k$ -regular graphs,  $k \geq 3$ .

A key ingredient (sometimes stated explicitly and often implicit in the proofs) in all the existing proofs of lower bounds for the Euclidean distortion of graphs is a Poincaré type inequality. Let  $G$  be a graph, and take any function  $f : V(G) \rightarrow \ell_2$ . A Poincaré type inequality bounds the average size of  $\{\|f(u) - f(v)\|_2\}_{u,v \in V(G)}$  in terms of its average "gradient"  $\{\|f(u) - f(v)\|\}_{[u,v] \in E(G)}$ . Such an inequality measures how often an equality holds in the triangle inequality. A lower bound on the Euclidean distortion of an  $n$ -point  $k$ -regular expander with conductance  $\Phi$ ,  $G$ , can be derived from the following Poincaré inequality for functions  $f : V(G) \rightarrow \mathbb{R}$  (see [M1]):

$$\sum_{u,v \in V(G)} |f(u) - f(v)| \leq \frac{n-1}{\Phi} \sum_{[u,v] \in E(G)} |f(u) - f(v)|.$$

For the embeddability of the Hamming cube  $D_n = \{0, 1\}^n$  in a metric space

$(X, d_X)$ , the following Poincaré inequality is relevant:

$$\sum_{d_H(u,v)=n} d_X^2(f(u), f(v)) \leq K^2 n^{\frac{2}{p}-1} \sum_{[u,v] \in E(D_n)} d_X^2(f(u), f(v)),$$

where  $d_H$  is the Hamming metric, and  $f : D_n \rightarrow X$  is any function. When such an inequality holds, for some  $p > 0$ , for every  $n$  and with  $K$  independent of  $n$  and  $f$ , we say that  $(X, d_X)$  has metric type  $p$  with constant  $K$ . This fundamental notion was introduced in [BouMW], where it was proved to essentially control the growth of  $c_X(D_n)$ . When  $X$  is a Banach space, metric type is studied via an important Poincaré inequality due to Pisier (See [P]. See also [W] for a different proof, and [T] for an improvement when  $X$  is the real line). We refer also to [BouMW], [P] for a related Poincaré inequality known as Enflo-type, or roundness.

Let  $T$  be the complete binary tree of depth  $N$ . The following Poincaré inequality is implicit in [LS], where a new proof of the estimate  $c_2(T_N) = \Omega(\sqrt{\log N})$  is obtained. Denote by  $r$  the root of  $T$ , and for every integer  $k$  let  $\mathcal{F}_k$  be the set of all unordered pairs of vertices  $\{u, v\} \subset V(T)$  such that  $d_T(u, r) = d_T(v, r)$  and  $d_T(u, v) = 2^k$ . Then, for every  $f : V(T) \rightarrow \mathbb{R}$ ,

$$\begin{aligned} \sum_{k=1}^{\lfloor \log_2 N \rfloor} \sum_{\{u,v\} \in \mathcal{F}_k} 2^{-d_T(u,r)-2^{k-1}-2k} \cdot |f(u) - f(v)|^2 \\ \leq C \sum_{[u,v] \in E(T)} 2^{-d_T(u,r)} |f(u) - f(v)|^2, \end{aligned}$$

where  $C$  is an absolute constant.

This paper further develops the above theme. We introduce two Poincaré type inequalities which are useful in the search of lower bounds for Euclidean distortion of graphs with large girth. The first is the notion of Markov type, due to K. Ball [B], which concerns the wandering of symmetric Markov chains whose state set is a metric space. We refer to section 2 for the definition. We also prove the following theorem, which can be viewed as a new Poincaré type inequality :

**Theorem 1.1.** *Let  $H$  be a Hilbert space and  $G$  be a  $k$ -regular graph,  $k \geq 3$ , with girth  $g$ . Fix some  $1 < s < g/2$ . For every  $f : V(G) \rightarrow H$  the following inequality holds:*

$$\sum_{d_G(u,v)=s} \|f(u) - f(v)\|^2 \leq Cs(k-1)^{s-1} \cdot \sum_{[u,v] \in E(G)} \|f(u) - f(v)\|^2,$$

where  $C$  is an absolute constant.

If, in addition, the graph  $G$  has a spectral gap, we can prove a stronger inequality. This leads to a new simple proof of the tightness of Bourgain's embedding theorem:

**Theorem 1.2.** *Let  $H$  be a Hilbert space and  $G$  be a  $k$ -regular graph,  $k \geq 3$ , with girth  $g$  and spectral gap  $\epsilon > 0$ . Fix some  $1 < s < g/2$ . For every  $f : V(G) \rightarrow H$  the following inequality holds:*

$$\sum_{d_G(u,v)=s} \|f(u) - f(v)\|^2 \leq C(k-1)^s \cdot \frac{1 - e^{-C\epsilon s/k}}{\epsilon} \cdot \sum_{[u,v] \in E(G)} \|f(u) - f(v)\|^2,$$

where  $C$  is an absolute numerical constant.

We apply the above inequalities to prove our main result:

**Theorem 1.3.** *There is a universal constant  $C > 0$  such that  $c_2(G) > C\sqrt{g}$  for every  $k$ -regular ( $k > 2$ ) graph  $G$  with girth  $g$ . If, in addition,  $G$  has a spectral gap  $\epsilon > 0$  then*

$$c_2(G) \geq \frac{Cg}{\sqrt{\min\{g, \frac{k}{\epsilon}\}}}.$$

REMARK. It is well known and not hard to show by probabilistic arguments that for every integer  $k \geq 3$ , there is an  $\epsilon = \epsilon_k > 0$  and  $n_0 = n_0(k)$  such that if  $n \geq n_0$  and  $kn$  is even, there exist  $k$ -regular graphs of order  $n$ , spectral gap greater than  $\epsilon$  and girth  $\Omega(\log n)$ . In view of Theorem 1.3 these graphs show that Bourgain's upper bound is tight. A curious feature of our estimate is that the lower bound for the Euclidean distortion of constant degree graphs with girth  $g$  does not depend on the graph's degree.

This paper contains two proofs for the first part of Theorem 1.3. We first present a proof based on the notion of Markov type. Next, we prove the inequalities in Theorems 1.1 and 1.2 to deduce the full statement of Theorem 1.3. This proof is based on quadratic programming. In section 4 we discuss the interrelations between the two methods.

## 2 A Proof Based on the Concept of Markov Type

The first proof we present is based on the important notion of Markov type, due to Ball [B]. This concept is a Lipschitz invariant of metric spaces. It is related to other "types" that are central to the modern theory of Banach spaces. The basic assumption of this concept can also be viewed as a Poincaré inequality on metric spaces. Although we will see later that the Markov type method cannot yield the second statement in Theorem 1.3,

it does give a simple and conceptual proof of the first statement. We first recall some definitions from [B].

Let  $(X, d)$  be a metric space. A symmetric Markov chain on  $X$  is a Markov chain  $\{Z_k\}_{k=0}^\infty$  on a state space  $\{x_1, \dots, x_m\} \subset X$  with a symmetric transition matrix and such that  $Z_0$  is uniformly distributed on  $\{x_1, \dots, x_m\}$ . In other words, there is a  $m \times m$  symmetric stochastic matrix  $A = (a_{ij})$  such that for all  $k$ ,  $P(Z_{k+1} = x_j | Z_k = x_i) = a_{ij}$  and  $P(Z_0 = x_i) = 1/m$ . For  $p > 0$  and an integer  $T$  let  $M_p(X, T)$  be the smallest constant  $C > 0$  such that for every symmetric Markov chain on  $X$ ,  $\{Z_k\}_{k=1}^\infty$

$$\mathbb{E} d^p(Z_T, Z_0) \leq C^p T \mathbb{E} d^p(Z_1, Z_0).$$

We say that  $(X, d)$  has Markov type  $p$  if  $M_p(X) := \sup_T M_p(X, T) < \infty$ . In this case  $M_p(X)$  is called the Markov type  $p$  constant of  $X$ .

The space  $L_2$  has Markov type 2 with constant 1, as shown in [B]. For the sake of completeness, we prove here a somewhat stronger and rather intuitive claim. We first observe that the Markov type 2 property for  $\mathbb{R}$  implies, by integration, the same conclusion for  $L_2$ . For symmetric Markov chains on  $\mathbb{R}$  we prove the following negative correlation inequality that implies Markov type 2. Let  $\{Z_k\}_{k=1}^\infty$  be a symmetric Markov chain with transition matrix  $A$  and state space  $\{x_1, \dots, x_m\} \subset \mathbb{R}$ . The symmetry assumption makes it intuitively plausible that  $Z_T - Z_{T-1}$  and  $Z_{T-1} - Z_0$  must be negatively correlated. To prove this, notice that  $Z_k$  is uniformly distributed on  $\{x_1, \dots, x_m\}$ , for every  $k$ . Since  $A$  is symmetric and stochastic, its spectrum is in  $[-1, 1]$ , and we deduce that  $(I - A)(I - A^k)$  is positive semi-definite for every  $k$ . Therefore,

$$\begin{aligned} \mathbb{E}(Z_T - Z_{T-1})(Z_{T-1} - Z_0) &= \mathbb{E}Z_T Z_{T-1} - \mathbb{E}Z_T Z_0 - \mathbb{E}Z_{T-1}^2 + \mathbb{E}Z_{T-1} Z_0 \\ &= \frac{1}{m} \left[ \sum_{i,j=1}^m (A)_{ij} x_i x_j - \sum_{i,j=1}^m (A^T)_{ij} x_i x_j - \sum_{i=1}^m x_i^2 + \sum_{i,j=1}^m (A^{T-1})_{ij} x_i x_j \right] \\ &= -\frac{1}{m} \langle (I + A^T - A - A^{T-1})x, x \rangle = -\frac{1}{m} \langle (I - A)(I - A^{T-1})x, x \rangle \leq 0, \end{aligned}$$

where  $x = (x_1, \dots, x_m)$ . Hence,

$$\begin{aligned} \mathbb{E}(Z_T - Z_0)^2 &= \mathbb{E}(Z_{T-1} - Z_0 + Z_T - Z_{T-1})^2 \\ &= \mathbb{E}(Z_{T-1} - Z_0)^2 + 2\mathbb{E}(Z_T - Z_{T-1})(Z_{T-1} - Z_0) + \mathbb{E}(Z_T - Z_{T-1})^2 \\ &\leq \mathbb{E}(Z_{T-1} - Z_0)^2 + \mathbb{E}(Z_1 - Z_0)^2. \end{aligned}$$

By summing this inequality over  $T = 1, \dots, N$ , we deduce that the real line has Markov type 2 with constant 1.

The following simple consequence of the above analysis will be useful for us:

COROLLARY 2.1. For every metric space  $(X, d)$ ,  $c_2(X, d) \geq M_2(X)$ .

*Proof.* Fix some embedding  $f : X \rightarrow L_2$  such that  $1/D \leq \|f(x) - f(y)\|/d(x, y) \leq 1$  for every  $x, y \in X$ . For every symmetric Markov chain  $\{Z_k\}_{k=0}^\infty$  on  $X$ , the Markov type 2 property of  $L_2$  applied to the Markov chain  $\{f(Z_k)\}_{k=0}^\infty$  gives

$$\begin{aligned} \frac{1}{D^2} \mathbb{E} d^2(Z_T, Z_0) &\leq \mathbb{E} \|f(Z_T) - f(Z_0)\|^2 \leq T \mathbb{E} \|f(Z_1) - f(Z_0)\|^2 \\ &\leq T \mathbb{E} d^2(Z_1, Z_0), \end{aligned}$$

so that  $D \geq M_2(X)$ . □

The first assertion in Theorem 1.3 follows from the following:

PROPOSITION 2.2. Let  $G$  be a  $k$ -regular graph with girth  $g$ . Then

$$M_2(G) \geq \frac{k-2}{k} \sqrt{\left\lceil \frac{g}{2} - 1 \right\rceil}.$$

*Proof.* Consider the symmetric Markov chain  $\{Z_t\}_{t=0}^\infty$  that corresponds to the canonical random walk on  $G$ . Namely,  $Z_0$  is uniformly distributed on  $V(G)$  and  $P(Z_{t+1} = v | Z_t = u)$  equals  $1/k$  if  $u$  and  $v$  are neighbors, and 0 otherwise. Note that every vertex  $v \in V(G)$  is the root of an induced  $k$ -regular tree of depth  $g/2$ . In other words, the metric of each ball of radius smaller than  $g/2$  in  $G$  is isometric to such a tree, whose root is the center of the ball. As long as  $T < g/2$ , each step of the random walk  $\{Z_t\}_{t=0}^T$  moves away from  $Z_0$  with probability at least  $\frac{k-1}{k}$  (if  $Z_t = Z_0$  then this probability is 1) and towards it with probability at most  $1/k$ . In other words, as long as  $T < g/2$ , the random walk has a positive drift away from  $Z_0$ . To quantify this, for every  $1 < T < g/2$  we have

$$\begin{aligned} \mathbb{E} d_G(Z_T, Z_0) &\geq \frac{k-1}{k} (\mathbb{E} d_G(Z_{T-1}, Z_0) + 1) + \frac{1}{k} (\mathbb{E} d_G(Z_{T-1}, Z_0) - 1) \\ &= \mathbb{E} d_G(Z_{T-1}, Z_0) + \frac{k-2}{k}. \end{aligned}$$

Hence, for every  $T < g/2$ ,

$$\mathbb{E} d_G^2(Z_T, Z_0) \geq (\mathbb{E} d_G(Z_T, Z_0))^2 \geq \left(\frac{k-2}{k}\right)^2 T^2.$$

On the other hand,

$$\mathbb{E} d_G^2(Z_T, Z_0) \leq M_2(G)^2 T \mathbb{E} d_G^2(Z_1, Z_0) = M_2(G)^2 T.$$

The proposition follows by taking  $T = \lceil \frac{g}{2} - 1 \rceil$ . □

### 3 Bounding the Distortion via Poincaré Inequalities

In what follows  $G$  is a  $k$  regular graph with girth  $g$ .

Semi-definite programming has proved to be a central tool in establishing lower bounds on  $c_2(\cdot)$ . Our proof of Theorems 1.1 and 1.2, and our second proof of Theorem 1.3 are all based on this point of view, together with an analysis of the algebraic properties of the graphs in question. We first present the necessary background.

Let  $PSD_n$  be the cone of positive semi-definite symmetric  $n \times n$  matrices. Define  $\mathcal{B}_n$  to be  $\{Q \in PSD_n \mid Q\mathbf{1} = 0\}$ , and let

$$\delta(Q, d) = \left( \frac{\sum_{i,j:Q_{i,j}>0} d^2(i,j)Q_{i,j}}{\sum_{i,j:Q_{i,j}<0} d^2(i,j)|Q_{i,j}|} \right)^{1/2},$$

if the denominator is not 0, and 1 otherwise.

The following lemma gives a formula for  $c_2$ .

**PROPOSITION 3.1** (Linial, London, Rabinovich [LLR]). *Suppose  $X$  is finite,  $|X| = n$ , then*

$$c_2(X, d) = \sup_{Q \in \mathcal{B}_n} \delta(Q, d)$$

In order to make use of the algebraic properties of  $G$ , we turn to some background on the following very useful concept.

**3.1 Geronimus polynomials.** Let  $G$  be a  $k$ -regular graph with girth  $g$  and let  $A$  be its adjacency matrix. We define  $A^{(t)}$  as  $G$ 's distance  $t$  matrix. Namely,  $A^{(t)}_{i,j} = 1$  if the distance  $d_G(i, j) = t$  and 0 otherwise.

There exist polynomials  $P_t$ , such that  $P_t$  has degree  $t$  and  $P_t(A) = A^{(t)}$  for every  $t < g/2$ . The conditions that define these polynomials easily translate to a simple recurrence relation. Clearly  $P_0(x) = 1$ , and  $P_1(x) = x$ . Note that  $A^{(t)} - A \cdot A^{(t-1)}$  equals  $-kA^{(t-2)} = -k \cdot I$  for  $t = 2$ , and  $-(k - 1)A^{(t-2)}$  for  $2 < t < g/2$ . Therefore,

$$P_2(x) = xP_1(x) - kP_0(x) = x^2 - k,$$

and

$$P_t(x) = xP_{t-1}(x) - (k - 1)P_{t-2}(x) \quad \text{for every } t > 2.$$

These polynomials are often called in the literature ‘‘Geronimus Polynomials’’, a name that we adopt. Basic facts about Geronimus polynomials can be found in [Bi], [S] (but note the different normalization used in these references). To make this discussion self-contained, we briefly review some of the necessary facts and sketch their proofs.

In order to understand the analytical properties of the Geronimus polynomials, one first solves the recursion and finds an explicit formula for

them. The following trigonometric expression is obtained:

$$\forall t > 0 \quad P_t(2\sqrt{k-1} \cos \theta) = (k-1)^{t/2-1} \frac{(k-1) \sin((t+1)\theta) - \sin((t-1)\theta)}{\sin \theta}. \tag{1}$$

To verify this identity, check the cases  $t = 1, 2$  and note that for  $t > 2$  the recursion relation holds.

Our next observation is that all the roots of  $P_t$  are real and they all lie between  $-2\sqrt{k-1}$  and  $2\sqrt{k-1}$ . This can be derived from the general theory of orthogonal polynomials (e.g. [Sz]), but we provide a direct proof. By identity 1 it suffices to show there are  $t$  distinct real values of  $\theta$  in  $[0, \pi)$  for which the above expression vanishes. Indeed, define  $\theta_q = (\frac{\pi}{2} + q\pi)/(t+1)$  for  $q = 0, 1, \dots, t-1$ . Now, it is not hard to see that  $P_t(2\sqrt{k-1} \cos \theta_q)$  is positive for even  $q$ , and negative for  $q$  odd. Therefore, there is a zero for some value of  $\theta$  between  $\theta_q$  and  $\theta_{q+1}$ , yielding  $t$  zeros in the desired interval.

The last two facts that we need are easily verified by induction:

$$P_t(k) = k(k-1)^{t-1} \quad \forall t > 0,$$

and

$$P'_t(k) = \frac{1}{(k-2)^2} (t(k-1)^{t+1} - 2(k-1)^t - t(k-1)^{t-1} + 2).$$

**3.2 Technical lemmas.** Even though the Geronimus polynomials are not convex throughout  $[-k, k]$ , we now prove an inequality that reflects the fact that their non-convexity is restricted to the relatively small range  $[-2\sqrt{k-1}, 2\sqrt{k-1}]$ . The proof uses the classical Markov inequality (see [BoE]):  $\|P'\|_{L_\infty[-1,1]} \leq n^2 \|P\|_{L_\infty[-1,1]}$  for every real polynomial  $P$  of degree  $n$ , where  $\|f\|_{L_\infty[-a,a]} = \sup_{|x| \leq a} |f(x)|$ . A more direct proof can be given by differentiating formula 1, but use of Markov's inequality eliminates a tedious calculation which leads, essentially, to the same estimate.

**LEMMA 3.2.** *Let  $s \geq 40$  be an even integer. For every  $\epsilon > 0$  and  $x \in [-k, k - \epsilon]$ ,*

$$\frac{P_s(k) - P_s(k - \epsilon)}{\epsilon} \geq \frac{P_s(k) - P_s(x)}{k - x}.$$

*Proof.* Define

$$f(x) = \frac{P_s(k) - P_s(x)}{k - x}.$$

We need to show that  $f$  is non-decreasing on  $[-k, k]$ . By taking a derivative of the right-hand side and expanding, this follows from the claim that for all  $x \in [-k, k]$ ,

$$h(x) := P_s(x) + (k - x)P'_s(x) \leq P_s(k).$$

Note that  $h(k) = P_s(k)$  and, since  $P_s$  is an even function (for  $s$  even), it follows that  $h(-k) = P_s(k) - 2kP'_s(k) < P_s(k)$  (since  $P'_s(k) > 0$ ). It is therefore enough to show that  $h(x_0) \leq P_s(k)$  whenever  $h'(x_0) = 0$ . Now,  $h'(x) = (k-x)P''_s(x)$ , so that the zeros of  $h'$  coincide with the zeros of  $P''_s$ . Since  $P_s$  has all its roots in the interval,  $[-2\sqrt{k-1}, 2\sqrt{k-1}]$ , the same holds for  $P''_s$ . It therefore suffices to show that  $h(x) \leq P_s(k)$  throughout the interval  $[-2\sqrt{k-1}, 2\sqrt{k-1}]$ . Every point in this interval has the form  $x = 2\sqrt{k-1} \cos \theta$  for some  $0 \leq \theta \leq \pi$ . Using the trigonometric expression (1) we get

$$P_s(x) = P_s(2 \cos \theta \sqrt{k-1}) = (k-1)^{s/2-1} \frac{(k-1) \sin((s+1)\theta) - \sin((s-1)\theta)}{\sin \theta}.$$

It is easily verified that  $\sin r\alpha \leq r \sin \alpha$  for  $\alpha \in [0, \pi)$  and  $r \geq 1$ . Therefore,

$$\|P_s\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]} \leq (k-1)^{s/2-1} ((k-1)(s+1) + (s-1)).$$

Markov's inequality implies that

$$\|P'_s\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]} \leq \frac{s^2}{2\sqrt{k-1}} \cdot \|P_s\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]},$$

so that

$$\begin{aligned} \|h\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]} &\leq \|P_s\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]} + 2k\|P'_s\|_{L_\infty[-2\sqrt{k-1}, 2\sqrt{k-1}]} \\ &\leq (k-1)^{s/2-1} ((k-1)(s+1) + (s-1)) \left(1 + \frac{ks^2}{\sqrt{k-1}}\right) \\ &\leq k(k-1)^{s-1} = P_s(k). \end{aligned}$$

It is not hard to verify the last inequality for every  $k \geq 3$  and  $s \geq 40$ . □

We need one more fact concerning Geronimus polynomials:

LEMMA 3.3. For every integer  $s$  and  $\epsilon \leq k/20$ ,

$$P_s(k - \epsilon) \geq k(k-1)^{s-1} \cdot e^{-150\epsilon s/k}.$$

*Proof.* Let  $y_1, \dots, y_s$  be the roots of  $P_s$ . By the mean value theorem there is some  $a \in (k - \epsilon, k)$  such that

$$\begin{aligned} \log \left[ \frac{P_s(k)}{P_s(k - \epsilon)} \right] &= \epsilon \frac{P'_s(a)}{P_s(a)} \\ &= \epsilon \cdot \sum_{i=1}^s \frac{1}{a - y_i} \leq \frac{\epsilon s}{k - \epsilon - 2\sqrt{k-1}} \leq \frac{150\epsilon s}{k}, \end{aligned}$$

where we have used the facts that  $y_1, \dots, y_n \in [-2\sqrt{k-1}, 2\sqrt{k-1}]$ ,  $\epsilon \leq k/20$  and  $k \geq 3$ . This yields the required result since  $P_s(k) = k(k-1)^{s-1}$ . □

COROLLARY 3.4. *Let  $s \geq 40$  be an even integer and  $0 < \epsilon \leq k$ . Then*

$$1 - \frac{P_s(k - \epsilon)}{P_s(k)} \leq C(1 - e^{-C\epsilon s/k}),$$

for some absolute constant  $C > 0$ .

*Proof.* When  $\epsilon \leq k/20$  this follows from Lemma 3.3. When  $\epsilon > k/20$  the right-hand side is bounded from below by an absolute constant. The left-hand side is at most 2, since the minimum of  $P_s$  is attained in the interval  $[-2\sqrt{k-1}, 2\sqrt{k-1}]$ , and the bound obtained in the proof of Lemma 3.2 implies in particular that  $P_s(x) \geq -P_s(k)$  for every  $x$  in this interval.  $\square$

**3.3 Proof of Theorems 1.1 and 1.2.** Let  $G$  be a  $k$ -regular graph with girth  $g$  and fix an integer  $1 < s < g/2$ . For every two vertices  $u, v \in V(G)$  such that  $d_G(u, v) = s$  there is a unique path of length  $s$  joining  $u$  and  $v$ . In other words, there is a unique set of vertices  $\{w_{u,v}(i)\}_{i=0}^s \subset V(G)$  such that  $w_{u,v}(0) = u$ ,  $w_{u,v}(s) = v$  and for every  $i \geq 1$ ,  $[w_{u,v}(i-1), w_{u,v}(i)] \in E(G)$ . Each edge  $e \in E(G)$  appears in exactly  $s(k-1)^{s-1}$  such paths. Hence, for every metric space  $(X, d_X)$  and every mapping  $f : V(G) \rightarrow X$

$$\begin{aligned} \sum_{d_G(u,v)=s} d_X^2(f(u), f(v)) &\leq \sum_{d_G(u,v)=s} \left( \sum_{i=1}^s d_X(f(w_{u,v}(i-1)), f(w_{u,v}(i))) \right)^2 \\ &\leq \sum_{d_G(u,v)=s} s \sum_{i=1}^s d_X^2(f(w_{u,v}(i-1)), f(w_{u,v}(i))) \\ &= s^2(k-1)^{s-1} \sum_{[u,v] \in E(G)} d_X^2(f(u), f(v)). \end{aligned}$$

It follows that the inequalities of Theorems 1.1 and 1.2 are trivially true for bounded  $s$ , so that we may clearly assume that  $s \geq 40$ . By a similar argument, we can also assume that  $s$  is even. In addition, in both statements it is clearly enough to assume that  $f$  maps  $G$  into  $H = \mathbb{R}$ .

We begin with the the proof of Theorem 1.2. Let  $G$  satisfy the conditions of Theorem 1.2 and consider the following matrix:

$$Q = \alpha I - A + \beta A^{(s)},$$

where  $40 \leq s < g/2$  is an even integer, and

$$\begin{aligned} \alpha &= k - \frac{\epsilon P_s(k)}{P_s(k) - P_s(k - \epsilon)}, \\ \beta &= \frac{\epsilon}{P_s(k) - P_s(k - \epsilon)}. \end{aligned}$$

We claim that  $Q \in \mathcal{B}_n$ . Now,  $Q$  is clearly symmetric. Also, since  $A\vec{1} = k\vec{1}$ ,  $A^{(s)}\vec{1} = P_s(k)\vec{1}$  so that  $Q\vec{1} = 0$ . Other than the eigenvalue  $k$ , the spectrum of  $A$  is in  $[-k, k - \epsilon]$ , so  $Q$  will be shown to be positive semi-definite if for all  $x \in [-k, k - \epsilon]$

$$\alpha - x + \beta P_s(x) \geq 0,$$

which is precisely the statement of Lemma 3.2.

Now, for every  $f : V(G) \rightarrow \mathbb{R}$  we get that

$$\begin{aligned} 0 &\leq \langle Qf, f \rangle \\ &= \alpha \sum_{u \in V(G)} f(u)^2 - \sum_{u,v \in V(G)} A_{uv} f(u)f(v) + \beta \sum_{u,v \in V(G)} A_{uv}^{(s)} f(u)f(v) \\ &= \frac{1}{2} \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2 - \frac{\beta}{2} \sum_{d_G(u,v)=s} |f(u) - f(v)|^2, \end{aligned}$$

where in the last equality we have used the fact that  $\alpha - k + \beta P_s(k) = 0$ . Hence,

$$\begin{aligned} \sum_{d_G(u,v)=s} |f(u) - f(v)|^2 &\leq \frac{1}{\beta} \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2 \\ &= \frac{k(k-1)^{s-1}}{\epsilon} \left[ 1 - \frac{P_s(k-\epsilon)}{P_s(k)} \right] \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2 \\ &\leq C(k-1)^s \cdot \frac{1 - e^{-C\epsilon s/k}}{\epsilon} \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2, \end{aligned}$$

where we have used Corollary 3.4.

The proof of Theorem 1.1 runs along the same lines. We return to the above construction and let  $\epsilon \rightarrow 0$ . This yields a matrix:

$$\tilde{Q} = \left[ k - \frac{P_s(k)}{P'_s(k)} \right] I - A + \frac{1}{P'_s(k)} A^{(s)},$$

which by continuity is in  $\mathcal{B}_n$  as well. Arguing as above, we get the following inequality:

$$\begin{aligned} \sum_{d_G(u,v)=s} |f(u) - f(v)|^2 &\leq P'_s(k) \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2 \\ &= \frac{s(k-1)^{s+1} - 2(k-1)^s - s(k-1)^{s-1} + 2}{(k-2)^2} \sum_{[u,v] \in E(G)} |f(u) - f(v)|^2, \end{aligned}$$

which implies the required result. □

**3.4 Proof of Theorem 1.3.** We now deduce Theorem 1.3 from Theorems 1.1 and 1.2. Let  $G$  be a  $k$ -regular graph ( $k \geq 3$ ) with girth  $g$ . Take any embedding  $f : V(G) \rightarrow \ell_2$  such that for every  $u, v \in V(G)$

$$\frac{1}{D} \leq \frac{\|f(u) - f(v)\|}{d_G(u, v)} \leq 1.$$

Set  $s = \lfloor g/2 \rfloor - 1$ . Since there are  $k(k - 1)^{s-1}$  vertices of distance  $s$  from a fixed vertex we get that:

$$\sum_{d_G(u,v)=s} \|f(u) - f(v)\|^2 \geq \frac{s^2 k(k - 1)^{s-1} |V(G)|}{D^2},$$

and

$$\sum_{[u,v] \in E(G)} \|f(u) - f(v)\|^2 \leq k|V(G)|.$$

Theorem 1.1 now gives that

$$\frac{s^2 k(k - 1)^{s-1} |V(G)|}{D^2} \leq C s(k - 1)^s |V(G)|,$$

so that  $c_2(G) \geq c' \sqrt{g}$ . If in addition we assume that  $G$  has a spectral gap  $\epsilon$ , this reasoning gives

$$c_2(G) \geq c' g \sqrt{\frac{\epsilon/k}{1 - e^{-Cg\epsilon/k}}} \geq \frac{c'' g}{\sqrt{\min \{g, \frac{k}{\epsilon}\}}},$$

which finishes the proof of Theorem 1.3.

### 4 The Relationship Between the Two Methods

The proof of the Markov type 2 property of Hilbert space in [B], sheds some light on the connection between the two methods. In Proposition 3.1 we seek the matrix  $Q$  with maximal  $\delta(Q, d)$  over all  $Q \in \mathcal{B}_n$ . We will show how to view Corollary 2.1 as a restriction of Proposition 3.1, in the sense that only a subset of the matrices in  $\mathcal{B}_n$  can be used for the lower bound.

Let  $B$  be the symmetric stochastic transition matrix defining the Markov chain, and let  $\gamma \in (0, 1)$ . Now define

$$R = \frac{2\gamma - 1}{\gamma} I - B + \frac{(1 - \gamma)^2}{\gamma} \sum_{l \geq 0} (\gamma B)^l.$$

The spectrum of  $B$  clearly determines that of  $R$ : If  $\lambda$  is an eigenvalue of  $B$ , then  $\frac{2\gamma - 1}{\gamma} - \lambda + \frac{(1 - \gamma)^2}{\gamma(1 - \gamma\lambda)}$  is an eigenvalue of  $R$ . This expression is nonnegative for  $\lambda \in [-1, 1]$ , and 0 for  $\lambda = 1$ . This clearly means that

$R \in \mathcal{B}_n$ . If  $x_1, \dots, x_n$  are vectors in Hilbert space then,

$$\sum_{i,j} R_{i,j} \|x_i - x_j\|^2 \leq 0,$$

and so

$$(1 - \gamma) \sum_{ij} \left( \sum_l (\gamma B)^l \right)_{ij} \|x_i - x_j\|^2 \leq \gamma \sum_{ij} B_{ij} \|x_i - x_j\|^2.$$

As Ball shows by taking  $\gamma = 1 - \frac{1}{m}$ , this is equivalent to the fact that Hilbert space has Markov type 2. To be more precise, this is the original definition he gives to the metric property “having a Markov type 2”.

Thus the approach via Markov type can be viewed as a specialized version of the semidefinite programming method. This method is incomplete in that not every matrix in  $\mathcal{B}_n$  is attained from the above transformation of symmetric stochastic matrices. This method is still very useful, in that it allows one to draw on geometric and probabilistic intuitions. In contrast, the semi-definite approach greatly depends on successful clever guesses of the matrix  $Q$ .

The Poincaré inequality method can also be viewed as a restriction of the family  $\mathcal{B}_n$ . Indeed, if  $G$  is a graph and  $B$  is any matrix such that  $B_{uv} = 0$  whenever  $u$  and  $v$  are not neighbors in  $G$ , then for any matrix  $C$  and  $\alpha \in \mathbb{R}$ , the fact that the matrix  $\alpha I - B + C$  is in  $\mathcal{B}_n$  is equivalent to the Poincaré inequality:

$$\sum_{u,v \in V(G)} C_{uv} |f(u) - f(v)|^2 \leq \sum_{[u,v] \in E(G)} B_{uv} |f(u) - f(v)|^2,$$

for every  $f : V(G) \rightarrow \mathbb{R}$ . Thus, all the existing lower bounds for  $c_2(\cdot)$  are based on the study of the above subset of  $\mathcal{B}_n$ , which leads to geometrically intuitive, Poincaré inequality reasons for non-embeddability of graphs in Hilbert space. Hence, in a sense, the full strength of Proposition 3.1 is yet to be fully exploited.

The second part of Theorem 1.3 cannot be derived from Markov type considerations. Indeed, let  $G$  be any graph. If  $\{Z_k\}_{k=0}^\infty$  is a symmetric Markov chain on  $G$  such that  $P(Z_{k+1} = u | Z_k = u) = 0$  for every  $u \in V(G)$  then clearly  $\mathbb{E} d_G^2(Z_T, Z_0) \leq \text{diam}(G)^2 \mathbb{E} d_G^2(Z_1, Z_0)$  and by the triangle inequality  $\mathbb{E} d_G^2(Z_T, Z_0) \leq T^2 \mathbb{E} d_G^2(Z_1, Z_0)$ . Hence, for every  $T$ ,

$$M_2(G) \leq \min \left\{ \frac{\text{diam}(G)}{\sqrt{T}}, T \right\},$$

which implies that  $M_2(G) = O(\text{diam}(G)^{2/3})$ , while our second proof of Theorem 1.3 showed that there are graphs with  $c_2(G) = \Omega(\text{diam}(G))$ .

An interesting problem that still remains open is the determination of the worst possible behavior of  $c_2(G)$  over all  $k$ -regular,  $k \geq 3$ , graphs with girth  $g$ . Our present methods seem to break at  $\sqrt{g}$ . On the other hand if  $k$ -regular graphs,  $G$ , with girth  $g$  and  $c_2(G) = O(\sqrt{g})$  exist, they cannot be expanders in view of the known lower bound for the Euclidean distortion of expanders. This observation rules out most known constructions for graphs with large girth. Although examples of  $k$ -regular graphs with large girth which are not expanders are known, these specific examples seem highly complicated and far from Euclidean.

Another problem worth mentioning is whether the lower bound for  $c_2(G)$  still holds without the regularity assumptions, i.e. if we only assume that the graph has large girth and the degree of each vertex is greater than 2. Finally, it would be very interesting to find lower bounds for  $c_1(G)$  as a function of the girth of  $G$ . In particular, we do not know if  $c_1(G)$  must tend to infinity when the girth tends to infinity.

## References

- [B] K. BALL, Markov chains, Riesz transforms and Lipschitz maps, *Geometric And Functional Analysis* 2 (1992), 137–172.
- [Ba] Y. BARTAL, On approximating arbitrary metrics by tree metrics, in “STOC ’98 (Dallas, TX)”, Proc. of the 24th International Colloquium on Automata, Languages, and Programming, July 1997, ACM, New York (1998), 161–168.
- [Bi] N. BIGGS, *Algebraic Graph Theory* Cambridge University Press, Cambridge, second edition, 1993.
- [BoE] P. BORWEIN, T. ERDÉLYI, *Polynomials and Polynomial Inequalities*, Graduate Texts in Mathematics, 161. Springer-Verlag, New York, 1995.
- [Bou1] J. BOURGAIN, On Lipschitz embedding of finite metric spaces in Hilbert space, *Israel J. Math.* 52 (1985), 46–52.
- [Bou2] J. BOURGAIN, The metrical interpretation of superreflexivity in Banach spaces, *Israel J. Math.* 56 (1986), 222–230.
- [BouMW] J. BOURGAIN, V. MILMAN, H. WOLFSON, On type of metric spaces, *Trans. Amer. Math. Soc.* 294 (1986), 295–317.
- [E] P. ENFLO, On the nonexistence of uniform homeomorphisms between  $L_p$ -spaces, *Arkiv För Matematik*, 8 (1969), 103–105.
- [GNRS] A. GUPTA, I. NEWMAN, Y. RABINOVICH, A. SINCLAIR, Cuts, trees and  $\ell_1$ -embeddings of graphs, in “Symposium on Foundations of Computer Science” (1999), 399–409.
- [KPR] P. KLEIN, S.A. PLOTKIN, S.B. RAO, Excluded minors, network decomposition and multicommodity flow, in 25th Annual ACM Symposium on

- the Theory of Computing (1993), 682–690.
- [LLR] N. LINIAL, E. LONDON, YU. RABINOVICH, The geometry of graphs and some of its algorithmic applications, *Combinatorica* 15 (1995), 215–245.
- [LM] N. LINIAL, A. MAGEN, Least-distortion Euclidean embeddings of graphs: Products of cycles and expanders, *J. of Combinatorial Theory ser. B* 79 (2000), 157–171.
- [LS] N. LINIAL, M. SAKS, On the Euclidean distortion of complete binary trees, preprint.
- [M1] J. MATOUŠEK, Embedding expanders into  $\ell_p$ , *Israel J. Math.* 102 (1997), 189–197.
- [M2] J. MATOUŠEK, On embedding trees into uniformly convex Banach spaces, *Israel J. Math.* 114 (1999), 221–237.
- [P] G. PISIER, Probabilistic methods in the geometry of Banach spaces, *Probability and Analysis (Varena, Italy, 1985)*, Springer Lecture Notes in Math. 1206 (1986), 167–241.
- [R] S.B. RAO, Small distortion and volume preserving embeddings for planar and Euclidean metrics, in “15th Annual ACM Symposium on Computational Geometry” (1999), 300–306.
- [S] P. SOLÉ, The second eigenvalue of regular graphs of given girth, *J. Combinatorial Theory, Ser. B* 56 (1992), 239–249.
- [Sz] G. SZEGŐ, *Orthogonal Polynomials*. Fourth edition. American Mathematical Society, Colloquium Publications, Vol. XXIII. American Mathematical Society, Providence, R.I., 1975.
- [T] M. TALAGRAND, Isoperimetry, logarithmic Sobolev inequalities on the discrete cube, and Margulis’ graph connectivity theorem, *Geom. Funct. Anal.* 3 (1993), 295–314.
- [W] R. WAGNER, Notes on an inequality by Pisier for Functions on the Discrete Cube, *Geometric Aspects of Functional Analysis*, Springer Lecture Notes in Math. 1745 (2000), 263–268.

NATHAN LINIAL, Institute of Computer Science, Hebrew University, Jerusalem 91904, Israel  
`nati@cs.huji.ac.il`

AVNER MAGEN, Institute of Computer Science, Hebrew University, Jerusalem 91904, Israel  
`avnerma@cs.huji.ac.il`

ASSAF NAOR, Institute of Mathematics, Hebrew University, Jerusalem 91904, Israel  
`naor@math.huji.ac.il`

Submitted: July 2001

Revision: September 2001