



(This is a sample cover image for this issue. The actual cover is not yet available at this time.)

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Journal of Combinatorial Theory, Series A

www.elsevier.com/locate/jcta


Strong convergence in posets

 Amir Ban^a, Nati Linial^b
^a Center for the Study of Rationality, Hebrew University, Jerusalem, Israel

^b School of Computer Science and Engineering, Hebrew University, Jerusalem, Israel

ARTICLE INFO

Article history:

Received 26 July 2011

Available online xxxx

Keywords:

Poset

Strong convergence

Order

ABSTRACT

We consider the following solitaire game whose rules are reminiscent of the children's game of leapfrog. The game is played on a poset $(P, <)$ with n elements. The player is handed an arbitrary permutation $\pi = (x_1, x_2, \dots, x_n)$ of the elements in P . At each round an element may "skip over" a smaller element preceding it, i.e. swap positions with it. For example, if $x_i < x_{i+1}$, then it is allowed to move from π to the permutation $(x_1, x_2, \dots, x_{i-1}, x_{i+1}, x_i, x_{i+2}, \dots, x_n)$ of P 's elements. The player is to carry out such steps as long as such swaps are possible. When there are several consecutive pairs of elements that satisfy this condition, the player can choose which pair to swap next. Does the player's choice of swaps matter for the final permutation or is it uniquely determined by the initial order of P 's elements? The reader may guess correctly that the latter proposition is correct. What may be more surprising, perhaps, is that this question is not trivial. The proof works by constructing an appropriate system of invariants.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Let $(P, <)$ be a poset with n elements. We say that the elements $x, y \in P$ are **comparable** if either $x < y$ or $y < x$. Otherwise we say they are incomparable and write $x \parallel y$. Let $\pi = (x_1, x_2, \dots, x_n)$ be some permutation of P 's elements. A **swap** changes this permutation to $(x_1, x_2, \dots, x_{i-1}, x_{i+1}, x_i, x_{i+2}, \dots, x_n)$ for some index i . This swap is **permissible** if $x_i < x_{i+1}$. A permutation of P 's elements is called **terminal** if no swap is permissible. We say that π *converges* to a terminal permutation σ of P 's elements if it is possible to move from π to σ through some sequence of permissible swaps. Clearly, every sequence of permissible swaps is finite, since no two elements can be swapped

E-mail addresses: amirban@netvision.net.il (A. Ban), nati@cs.huji.ac.il (N. Linial).

more than once, and therefore every permutation converges. The main result in this note states that the convergence is *strong*, in the sense that the permutation converges to the same terminal permutation regardless of the order of swaps:

Theorem 1. *Every permutation π of the elements of a finite poset $(P, <)$ converges to exactly one terminal permutation.*

The question arose in the analysis of certain dynamic economic systems [2] in which agents compete against each other. An agent's standing in the process is quantified by a positive real number that captures his current *reputation*. As the process unfolds, reputations go up and down, but the absolute scores do not matter and the only issue is the agents' ranking by reputation. It can be shown that in the setting of [2] the possible order changes are governed by an (unknown, but fixed) partial order on the agents. However, the actual timing and dynamics at which agents move up and down in the reputation ranking is controlled by a complex set of stochastic rules. The question came up, then, what can be said about the players' ranking when the system reaches a steady state. As our result shows, the steady state of the dynamical system is determined solely by its initial state. We hope that the result may be useful in the study of dynamical systems in the physical and biological fields as well.

Elementary though this question is, it seems to be new. We note, however, that strong convergence has been studied before in combinatorics. Several beautiful theorems establish strong convergence phenomena in various "reflection games". The common theme is this: A graph $G = (V, E)$ is given along with a function $w : V \rightarrow \mathbb{Z}$. In addition, there is a rule that allows certain local modifications to w . Such local changes can take place only as long as w takes some negative values and the process terminates when $w \geq 0$. Several such games were analyzed in the literature [1,4,3,6,5]. All these papers show that even though there generally exist a number of possible allowed steps, every sequence of allowed steps terminates and the terminal position is independent of the steps chosen. In some cases it is even shown that the number of steps till termination depends only on the initial position. Despite the apparent similarity, we do not see a reduction between our results and these theorems.

2. The proof

Proof. Let π converge to some terminal permutation τ . The uniqueness of τ is proved by providing a criterion, depending only on π , as to which pairs of elements appear in the same order in π and τ and which are reversed.

An (x, y) -link in π is a sequence $x = z_1, z_2, \dots, z_k = y$ that appear in this order (not necessarily consecutively) in π such that $z_\alpha \parallel z_{\alpha+1}$ for every $\alpha \in [k - 1]$.

Clearly, if $x \parallel y$ no permissible swap can change the relative order of x and y . Consequently:

- No sequence of permissible swaps can change the relative order of x and y if an (x, y) -link exists.
- No sequence of permissible swaps can create or eliminate an (x, y) -link.

We say that (x, y) is a *critical pair* in π if (i) $x < y$, (ii) x precedes y in π and, (iii) there is no (x, y) -link in π .

We now assert and prove the criterion for whether any two elements x and y in π , with x preceding y , preserve or reverse their relative order in a terminal permutation:

1. If $y < x$, the order is preserved.
2. If there exists an (x, y) -link, the order is preserved.
3. Otherwise, i.e. if (x, y) is a critical pair, the order is reversed.

The first element of the criterion is trivial and the second has already been dealt with. It remains to show the third and last element: Since an (x, y) -link cannot be created or eliminated by permissible swaps, an equivalent statement to this claim is that a permutation τ with a critical pair cannot be terminal. We prove this by induction on the number of elements in τ separating x and y :

At the base of induction, if x and y are neighbor elements, the assertion is true since as $x < y$ the permutation is not terminal. Now let k be the number of elements separating x and y , and the induction hypothesis is that if the number of elements separating a pair is less than k it cannot be critical.

Let z be an element between x and y in τ . Assume $x < z$. Then by the induction hypothesis there exists an (x, z) -link. Now consider the relation between z and y : $z \parallel y$ is impossible, because then y could be concatenated to the (x, z) -link to form an (x, y) -link, contrary to the assumption that (x, y) is a critical pair. Similarly, $z < y$ would by the induction hypothesis prove the existence of a (z, y) -link, but this is impossible as it could be concatenated to the (x, z) -link to form an (x, y) -link. This leaves $y < z$ as the only possibility.

In summary $x < z \Rightarrow y < z$.

By similar reasoning $z < y \Rightarrow z < x$.

Furthermore, the possibility $x \parallel z$ together with $z \parallel y$ can be dismissed as constituting an (x, y) -link, leaving just two possible scenarios satisfied by each z between x and y :

- z is “small”, i.e. $z < x$ and $z < y$.
- z is “large”, i.e. $x < z$ and $y < z$.

Since τ is terminal, x 's immediate neighbor must be “small”, and y 's immediate neighbor must be “large”. Between these two, there must exist two consecutive elements z_1, z_2 such that z_1 is “small” and z_2 is “large”. But this leads to a contradiction as $z_1 < x < z_2 \Rightarrow z_1 < z_2$, which implies that τ is not terminal. Therefore (x, y) cannot be critical, completing the proof by induction that a terminal permutation cannot have a critical pair.

This completes the proof for the uniqueness of the terminal permutation. \square

3. Remarks

As the proof shows, the relative final order of every pair of elements x and y is determined by the existence of a link that connects them. If a link exists, then x, y maintain their initial relative order in the final permutation. If no such link exists, their final relative order agrees with their mutual order relation.

The proof also demonstrates that the number of swaps to reach termination is uniquely defined: Between the initial and terminal permutation, sum the displacements of elements that moved forward (alternatively sum the displacements of backward moving elements).

The result shows as well how to efficiently determine the terminal permutation given how the elements are ordered initially: Perform an arbitrary swap until no more swaps are possible.

One of the referees asked us whether the result extends from posets to directed acyclic graphs. At least at its simplest form the analogous claim is incorrect. Let D be the directed path on 3 vertices $A \rightarrow B \rightarrow C$. The permutation ABC converges to BAC if A, B are swapped, but to ACB if B, C are swapped.

It is of interest to understand how many terminal permutations various posets have. In particular how is this parameter distributed over all n -element posets?

References

- [1] N. Alon, I. Krasikov, Y. Peres, Reflection sequences, Amer. Math. Monthly 96 (1989) 820–822.
- [2] A. Ban, N. Linial, The dynamics of reputation systems, in: Proceedings of the 13th Conference on Theoretical Aspects of Rationality and Knowledge, 2011, pp. 91–100.
- [3] K. Eriksson, Strong convergence and a game of numbers, European J. Combin. 17 (4) (1996) 379–390.
- [4] S. Mozes, Reflection processes on graphs and Weyl groups, J. Combin. Theory Ser. A 53 (1990) 128–142.
- [5] R.P. Stanley, Enumerative Combinatorics, Cambridge University Press, Cambridge, 1999.
- [6] P. Winkler, Mathematical Puzzles: A Connoisseur's Collection, A. K. Peters, 2004.